arXiv:0910.4885v1 [math.RT] 26 Oct 2009

LOW-DIMENSIONAL COMPLEX CHARACTERS OF THE SYMPLECTIC AND ORTHOGONAL GROUPS

HUNG NGOC NGUYEN

Abstract. We classify the irreducible complex characters of the symplectic groups $Sp_{2n}(q)$ and the orthogonal groups $Spin^{\pm}_{2n}(q)$, $Spin_{2n+1}(q)$ of degrees up to the bound D, where $D = (q^n - 1)q^{4n-10}/2$ for symplectic groups, $D = q^{4n-8}$ for orthogonal groups in odd dimension, and $D = q^{4n-10}$ for orthogonal groups in even dimension.

1. Introduction

Lower bounds for the degrees of non-trivial irreducible representations of finite groups of Lie type in cross characteristic were found by Landazuri, Seitz, and Zalesskii in [LS], [SZ]. These bounds have proved to be very useful in various applications. Thanks to the effort of many people, the exact value of the smallest degree of non-trivial irreducible representations was determined for many groups of Lie type (see [T]).

We are interested in not only the smallest representation, but more importantly, the low-dimensional representations. In the case of complex representations, this problem was first studied by Tiep and Zalesskii [TZ1] for finite classical groups and then by Lübeck [Lu] for exceptional groups. For representations over fields of cross characteristics, this problem has been studied recently in [BK], [GT1] for $SL_n(q)$; [GMST], [HM] for $SU_n(q)$; [GMST] for $Sp_{2n}(q)$ with q odd; and [GT2] for $Sp_n(q)$ with q even. Let H be one of these groups and denote by $\mathfrak{d}(H)$ the smallest degree of nontrivial irreducible representations in cross characteristic. The purpose of these papers is to classify the irreducible representations of H of degrees close to $\mathfrak{d}(H)$ and to prove that there is a relatively "big" gap between the degrees of these representations and the next degree.

From now on, we denote by G either the symplectic groups $Sp_{2n}(q)$ or the orthogonal groups $Spin^{\pm}_{n}(q)$, where q is a power of a prime p. The smallest non-trivial

---

Date: October 26, 2009.

Part of this work was done during my PhD study at the University of Florida. I am grateful to my advisor, Professor P. H. Tiep, for the devoted guidance and fruitful discussion. I would like to thank Professor M. W. Liebeck for motivating this work. Also, I want to thank Professor J. I. Hall and Professor U. Meierfrankenfeld for their generous hospitality and support during my time at the Michigan State University. Finally, I thank the referee for his careful reading of the manuscript and helpful suggestions.

complex characters of G was determined in [TZ1]. It was also proved in [TZ1] that, up to the bound $(q^{2n}-1)/2(q+1)$, the odd characteristic symplectic group $Sp_{2n}(q)$ has four irreducible characters of degrees $(q^n \pm 1)/2$, which are so-called *Weil characters*, and the smallest unipotent character of degree $(q^n-1)(q^n-q)/2(q+1)$. We want to extend these results to a larger bound. More precisely, we classify the irreducible complex characters of G of degrees up to the bound D, where $D=(q^n-1)q^{4n-10}/2$ for the odd characteristic symplectic groups, $D=q^{4n-8}$ for the orthogonal groups in odd dimension, and $D=q^{4n-10}$ for the orthogonal groups in even dimension. When q is even, the irreducible complex characters of $Sp_{2n}(q) \simeq Spin_{2n+1}(q)$ were classified up to an already good enough bound $(q^{2n}-1)(q^{n-1}-1)(q^{n-1}-q^2)/2(q^4-1)$ in [GT2] and therefore it is not in our consideration. Theorems 1.2, 1.3, 1.4, and Proposition 8.3 in this paper have been used in [LOST] to establish the Ore's conjecture for the orthogonal groups. Our main results are the following theorems.

Theorem 1.1. *Let χ be an irreducible complex character of $G=Sp_{2n}(q)$, where $n \geq 6$ and q is an odd prime power. Then either $\chi(1) > (q^n-1)q^{4n-10}/2$ or χ belongs to a list of $q^2+12q+36$ characters in Tables 1, 2, 3 (at the end of §4).*

Theorem 1.2. *Let χ be an irreducible complex character of $G=Spin_{2n+1}(q)$, where $n \geq 5$ and q is an odd prime power. Then $\chi(1)=1$ (1 character), $\chi(1)=(q^{2n}-1)/(q^2-1)$ (1 character), $\chi(1)=q(q^{2n}-1)/(q^2-1)$ (1 character), $(q^n+\alpha_1)(q^n+\alpha_2 q)/2(q+\alpha_1\alpha_2)$ (4 characters, $\alpha_{1,2}=\pm 1$), $\chi(1)=(q^{2n}-1)/(q+\alpha)$ ($(q+\alpha-2)/2$ characters for each $\alpha=\pm 1$), or $\chi(1) > q^{4n-8}$.*

Theorem 1.3. *Let χ be an irreducible complex character of $G=Spin^{\alpha}_{2n}(q) \simeq \Omega^{\alpha}_{2n}(q)$, where $n \geq 5, \alpha = \pm$, $(n,q,\alpha) \neq (5,2,+)$, and q is a power of 2. Let*

$$D(n,q,\alpha)=\begin{cases} (q-1)(q^2+1)(q^3-1)(q^4+1), & n=5, \alpha=-, \\ q^{4n-10}+1, & \textit{otherwise}. \end{cases}$$

Then $\chi(1)=1$ (1 character), $\chi(1)=(q^n-\alpha)(q^{n-1}+\alpha q)/(q^2-1)$ (1 character), $\chi(1)=(q^{2n}-q^2)/(q^2-1)$ (1 character), $\chi(1)=(q^n-\alpha)(q^{n-1}+\alpha\beta)/(q-\beta)$ ($(q-\beta-1)/2$ characters for each $\beta=\pm$), or $\chi(1) \geq D(n,q,\alpha)$.

Furthermore, when $(n,\alpha)=(5,-)$, G has exactly q characters of degree $D(5,q,-)$, 1 character of degree $q^2(q^4+1)(q^5+1)/(q+1)$, and no more characters of degrees up to q^{10}.

Theorem 1.4. *Let χ be an irreducible complex character of $G=Spin^{\alpha}_{2n}(q)$, where $\alpha=\pm$, $n \geq 5$, and q is an odd prime power. Let*

$$D(n,q,\alpha)=\begin{cases} (q-1)(q^2+1)(q^3-1)(q^4+1), & n=5, \alpha=-, \\ q^{4n-10}+1, & \textit{otherwise}. \end{cases}$$

Then $\chi(1)=1$ (1 character), $\chi(1)=(q^n-\alpha)(q^{n-1}+\alpha q)/(q^2-1)$ (1 character), $\chi(1)=(q^{2n}-q^2)/(q^2-1)$ (1 character), $\chi(1)=(q^n-\alpha)(q^{n-1}-\alpha)/2(q+1)$ (2 characters),

$\chi(1) = (q^n - \alpha)(q^{n-1} + \alpha)/2(q-1)$ (2 characters), $\chi(1) = (q^n - \alpha)(q^{n-1} + \alpha\beta)/(q-\beta)$ ($(q-\beta-2)/2$ characters for each $\beta = \pm$), or $\chi(1) \geq D(n,q,\alpha)$.

Furthermore, when $(n,\alpha) = (5,-)$, G has exactly q characters of degree $D(5,q,-)$, 1 character of degree $q^2(q^4+1)(q^5+1)/(q+1)$, and no more characters of degrees up to q^{10}.

This paper is organized as follows. In §2, we recall the Lusztig's classification of the irreducible complex characters of finite groups of Lie type. Also, the structures of centralizers of semi-simple elements in symplectic and orthogonal groups are collected in §2. The low-dimensional unipotent characters of G are classified in §3. Each of the groups $Sp_{2n}(q)$ (q odd), $Spin_{2n+1}(q)$ (q odd), $Spin^{\pm}_{2n}(q)$ (q even), and $Spin^{\pm}_{2n}(q)$ (q odd) is treated individually in §4, §5, §6, and §7, respectively. Finally, in §8, we count the number of irreducible complex characters of $G = Spin^{\pm}_{12}(3)$ of degrees up to $4 \cdot 3^{15}$.

2. Preliminaries

2.1. **Notation.** Throughout this paper, $GL^+_n(q)$ stands for $GL_n(q)$ and $GL^-_n(q)$ stands for $GU_n(q)$. Furthermore, ± could be understood as ± itself or ±1, depending on the context. In order to label the eigenvalues of semi-simple elements, as well as conjugacy classes of finite classical groups, we follow the notation of [S] and [W2]. Let κ denote a generator of the field of q^4 elements, $\zeta = \kappa^{q^2-1}$, $\theta = \kappa^{q^2+1}$, $\eta = \theta^{q-1}$, $\gamma = \theta^{q+1}$. Let $T_1 = \{1, ..., (q-3)/2\}$, $T_2 = \{1, ..., (q-1)/2\}$ if q is odd and $T_1 = \{1, ..., (q-2)/2\}$, $T_2 = \{1, ..., q/2\}$ if q is even. Furthermore, when q is odd, let $R_1^* = \{j \in \mathbb{Z} : 1 \leq j < q^2+1, j \neq (q^2+1)/2\}$ and define R_1 to be a complete set of class representatives of the equivalence relation $\sim$ on R_1^*:

$$i \sim j \Leftrightarrow i \equiv \pm j \text{ or } \pm qj \pmod{q^2+1}.$$

Similarly, let $R_2^* = \{j \in \mathbb{Z} : 1 \leq j \leq q^2-1, q-1 \nmid j, q+1 \nmid j\}$ and define R_2 to be a complete set of class representatives of the equivalence relation $\sim$ on R_2^*:

$$i \sim j \Leftrightarrow i \equiv \pm j \text{ or } \pm qj \pmod{q^2-1}.$$

Note that $|R_1| = (q^2-1)/4$ and $|R_2| = (q-1)^2/4$.

2.2. **Lusztig's classification.** The main ingredient to prove our results is Lusztig's classification of irreducible complex characters of finite groups of Lie type. To make the paper self-contained, we recall it here.

Let G be either the symplectic groups $Sp_{2n}(q)$ or the orthogonal groups $Spin^{\pm}_{2n}(q)$, $Spin_{2n+1}(q)$ where q is a power of a prime p. Let $\mathcal{G}$ be the algebraic group and F the Frobenius endomorphism on $\mathcal{G}$ such that $G = \mathcal{G}^F$. Let $\mathcal{G}^*$ be the dual group of $\mathcal{G}$ and F^* the dual Frobenius endomorphism and denote $G^* = \mathcal{G}^{*F^*}$. Lusztig's classification (see chapter 13 of [DM]) says that the set of irreducible complex characters of G is partitioned into Lusztig series $\mathcal{E}(G,(s))$ associated to various geometric conjugacy classes (s) of semi-simple elements of G^*. In fact, $\mathcal{E}(G,(s))$ is the set of irreducible constituents of a Deligne-Lusztig character $R^{\mathcal{G}}_{\mathcal{T}}(\theta)$, where $(\mathcal{T},\theta)$ is of the geometric conjugacy class associated to (s). The elements of $\mathcal{E}(G,(1))$ are called *unipotent characters* of G. When $\mathcal{G}$ is a connected reductive group, for any semi-simple element $s \in G^*$, there is a bijection $\chi \mapsto \psi$ from $\mathcal{E}(G,(s))$ to $\mathcal{E}(C_{G^*}(s),(1))$ such that

$$\chi(1) = \frac{|G|_{p'}}{|C_{G^*}(s)|_{p'}}\psi(1). \tag{2.1}$$

In this situation, we will say that the irreducible complex character χ of G is parameterized by the pair $((s),\psi)$.

Suppose that we are determining irreducible complex characters of G of degrees up to a certain bound D. Unipotent characters of G as well as G^* are classified by Lusztig (see §13.8 of [C1]) and we will use that to find the low-dimensional unipotent characters of G. When the character χ is not unipotent, i.e. $(s) \neq (1)$, based on the structure of $C_{G^*}(s)$, we estimate $(G^* : C_{G^*}(s))_{p'}$ and come up with certain cases when $C_{G^*}(s)$ is large enough. More specifically, from formula 2.1, we see that if $\chi(1) < D$ then $|C_{G^*}(s)| > |G|_{p'}/D$. The following Proposition is used frequently to determine unipotent characters of $C_{G^*}(s)$.

Proposition 2.1 (Proposition 13.20 of [DM]). *Let $\mathcal{G}$ and $\mathcal{G}_1$ be two reductive groups defined over $\mathbb{F}_q$, and let $f : \mathcal{G} \to \mathcal{G}_1$ be a morphism of algebraic groups with a central kernel, defined over $\mathbb{F}_q$ and such that $f(\mathcal{G})$ contains the derived group $\mathcal{G}_1'$; then the unipotent characters of $\mathcal{G}^F$ are the $\theta \circ f$, where θ runs over the unipotent characters of $\mathcal{G}_1^F$.*

2.3. **Centralizers of semi-simple elements.** We collect here some well-known results about the structures of centralizers of semi-simple elements in finite classical groups (cf. [C2], [FS], and [TZ2]). Since the following lemmas are similar, we omit most of their proofs except the last one's.

Lemma 2.2. *Let q be odd and $s \in SO_{2n+1}(q)$ be semi-simple. Then*

$$C_{SO_{2n+1}(q)}(s) \simeq SO_{2k+1}(q) \times O^{\pm}_{2(m-k)}(q) \times \prod_{i=1}^{t} GL^{\alpha_i}_{a_i}(q^{k_i})$$

and

$$C_{O_{2n+1}(q)}(s) \simeq O_{2k+1}(q) \times O^{\pm}_{2(m-k)}(q) \times \prod_{i=1}^{t} GL^{\alpha_i}_{a_i}(q^{k_i}),$$

where $0 \leq k \leq m \leq n$, $\alpha_i = \pm$, and $\sum_{i=1}^{t} k_i a_i = n-m$.

Lemma 2.3. *Let q be even and $s \in O^{\pm}_{2n}(q)$ be semi-simple. Then*

$$C_{O^{\pm}_{2n}(q)}(s) \simeq O^{\pm}_{2m}(q) \times \prod_{i=1}^{t} GL^{\alpha_i}_{a_i}(q^{k_i}),$$

where $\alpha_i = \pm$ *and* $\sum_{i=1}^{t} k_i a_i = n - m$.

Let $(.,.)$ be a non-degenerate symplectic form on the space $V = \mathbb{F}_q^{2n}$, then the *conformal symplectic group* $CSp_{2n}(q)$ is defined to be

$$\{g \in GL(V) \mid \exists\, \tau(g) \in \mathbb{F}_q^*, \forall u, v \in V, (gu, gv) = \tau(g)(u, v)\}.$$

Lemma 2.4. *Let q be odd and $s \in CSp_{2n}(q)$ be semi-simple. Then*

$$C_{CSp_{2n}(q)}(s) \simeq C_{Sp_{2n}(q)}(s) \cdot \mathbb{Z}_{q-1}.$$

Moreover,

(i) *If $\tau(s)$ is not a square in $\mathbb{F}_q$, then*

$$C_{Sp_{2n}(q)}(s) \simeq Sp_m(q^2) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i}),$$

where m is even, $\alpha_i = \pm$, and $\sum_{i=1}^{t} k_i a_i = n - m$.

(ii) *If $\tau(s)$ is a square in $\mathbb{F}_q$, then*

$$C_{Sp_{2n}(q)}(s) \simeq Sp_{2k}(q) \times Sp_{2(m-k)}(q) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i}),$$

where $0 \le k \le m \le n$, $\alpha_i = \pm$, and $\sum_{i=1}^{t} k_i a_i = n - m$.

Let $Q(.)$ be a non-degenerate quadratic form on the space $V = \mathbb{F}_q^{2n}$. Then the *conformal orthogonal group* $CO_{2n}^{\pm}(q)$ is defined to be

$$\{g \in GL(V) \mid \exists\, \tau(g) \in \mathbb{F}_q^*, \forall v \in V, Q(g(v)) = \tau(g)Q(v)\}.$$

We use the description of $CO_{2n}^{\pm}(q)^0$ in [TZ1, Remark 7.3]. When q is odd, it is also shown in [TZ1, Lemma 7.4] that $CO_{2n}^{\pm}(q)^0 = \{g \in CO_{2n}^{\pm}(q) \mid \det(g) = \tau(g)^n\}$ and $CO_{2n}^{\pm}(q)^0$ is actually a subgroup of index 2 of $CO_{2n}^{\pm}(q)$ such that $CO_{2n}^{\pm}(q)^0/SO_{2n}^{\pm}(q) \simeq CO_{2n}^{\pm}(q)/O_{2n}^{\pm}(q) \simeq \mathbb{Z}_{q-1}$.

Lemma 2.5. *Let q be odd and $s \in CO_{2n}^{\pm}(q)^0$ be semi-simple. Then*

$$C_{CO_{2n}^{\pm}(q)}(s) \simeq C_{O_{2n}^{\pm}(q)}(s) \cdot \mathbb{Z}_{q-1},$$

and

$$C_{CO_{2n}^{\pm}(q)^0}(s) \simeq C_{SO_{2n}^{\pm}(q)}(s) \cdot \mathbb{Z}_{q-1}.$$

Moreover,

(i) *If $\tau(s)$ is not a square in $\mathbb{F}_q$, then*

$$C_{O_{2n}^{\pm}(q)}(s) \simeq O_m^{\pm}(q^2) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i}),$$

where m is even, $\alpha_i = \pm$, and $\sum_{i=1}^{t} k_i a_i = n - m$.

(ii) *If* $\tau(s)$ *is a square in* $\mathbb{F}_q$, *then*

$$C_{O^{\pm}_{2n}(q)}(s) \simeq O^{\pm}_{2k}(q) \times O^{\pm}_{2(m-k)}(q) \times \prod_{i=1}^{t} GL^{\alpha_i}_{a_i}(q^{k_i}),$$

where $0 \le k \le m \le n$, $\alpha_i = \pm$, *and* $\sum_{i=1}^{t} k_i a_i = n - m$.

Proof. Let $(.,.)$ be the non-degenerate bilinear form associated with $Q(.)$. Fix a basis of V and let J be the Gram matrix of $(.,.)$ corresponding to this basis. Then ${}^t sJs = \tau J$, where $\tau := \tau(s)$. Hence $\mathrm{Spec}(s) = \mathrm{Spec}({}^t s) = \tau \mathrm{Spec}(Js^{-1}J^{-1}) = \tau \mathrm{Spec}(s^{-1})$. Denote the characteristic polynomial of s acting on V by $P(x) \in \mathbb{F}_q[x]$. There are two cases:

Case 1: If τ is not a square in $\mathbb{F}_q$, suppose $\tau = \lambda_1^2$ where $\lambda_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Decompose $P(x)$ into irreducible polynomials over $\mathbb{F}_q$:

$$P(x) = (x^2 - \tau)^m \prod_{i=1}^{l} f_i^{m_i}(x) \prod_{j=1}^{l'} g_j^{n_j}(x) \widehat{g_j}^{n_j}(x),$$

where

- If λ is a root of $f_i(x)$ then $\tau\lambda^{-1}$ is also a root of $f_i(x)$, $\deg(f_i)$ is even, $\pm\lambda_1$ are not the roots of f_i,
- If λ is a root of $g_j(x)$ then $\tau\lambda^{-1}$ is a root of $\widehat{g_j}(x)$, $\deg(g_j) = \deg(\widehat{g_j})$, $g_j \neq \widehat{g_j}$,
- $n = m + \Sigma_{i=1}^{l} m_i \deg(f_i)/2 + \Sigma_{j=1}^{l'} n_j \deg(g_j)$.

Since $s \in CO^{\pm}_{2n}(q)^0$, we have $\tau^n = \det(s) = (-\tau)^m \tau^{n-m}$. Therefore, m is even. Set

$$V_\tau := (\frac{P(x)}{(x^2-\tau)^m})_{x=s}(V), V_i := (\frac{P(x)}{f_i^{m_i}(x)})_{x=s}(V), U_j := (\frac{P(x)}{g_j^{n_j}(x)\widehat{g}_j^{n_j}(x)})_{x=s}(V).$$

Then $V = V_\tau \oplus V_1 \oplus \cdots V_l \oplus U_1 \oplus \cdots \oplus U_{l'}$ is an orthogonal decomposition of V. Let $W \in \{V, V_\tau, V_1, ..., V_l, U_1, ..., U_{l'}\}$. Then $s|_W$ is semi-simple and $(.,.)$ is non-degenerate on W. For short notation, we denote $C_{O(W)}(s|_W)$ and $C_{CO(W)}(s|_W)$ by $C_{O(W)}(s)$ and $C_{CO(W)}(s)$, respectively.

1a) First, we show that $C_{O(W)}(s) \simeq GU_{m_0}(q^k)$ and $C_{CO(W)}(s) \simeq C_{O(W)}(s) \cdot \mathbb{Z}_{q-1}$, where $W = V_i$ for $i = 1, ..., l$, $m_0 = m_i$, and $k = \deg(f_i)/2$. Note that the characteristic polynomial of s acting on W is $f_i^{m_0}(x)$. Let $\lambda \in \mathbb{F}_{q^{2k}}$ be an eigenvalue of the action of s on W. Then all eigenvalues of the action of s on W are $\lambda, \lambda^q, ..., \lambda^{q^{2k-1}}$. Since $\tau\lambda^{-1}$ is also a root of $f_i(x)$ and $\pm\lambda_1$ are not, $\tau\lambda^{-1} = \lambda^{q^k}$. Consider $\widetilde{W} = W \otimes_{\mathbb{F}_q} \mathbb{F}_{q^{2k}}$. Fix a basis (f_i) in W, and define a Frobenius endomorphism $\sigma : \Sigma_i x_i f_i \mapsto \Sigma_i x_i^q f_i$ on $\widetilde{W}$, where $x_i \in \mathbb{F}_{q^{2k}}$. The simplicity of s implies that $\widetilde{W} = \widetilde{W_1} \oplus \cdots \oplus \widetilde{W_{2k}}$, where $\widetilde{W_i} = \mathrm{Ker}(s - \lambda^{q^{i-1}})$. We see that σ permutes the $\widetilde{W_i}$'s cyclically: $\sigma(\widetilde{W_i}) = \widetilde{W_{i+1}}$, where $\widetilde{W_{2k+1}} = \widetilde{W_1}$. Let $g \in CO(W)$ be commuting with $s|_W$. Then g preserves each $\widetilde{W_i}$. Moreover, it is easy to see that g also commutes with σ. This implies that the action of g on $\widetilde{W}$ is completely determined by its action on $\widetilde{W}_1$: $g(\sigma^i w) = \sigma^i(gw)$ for $w \in \widetilde{W}_1$. So $C_{CO(W)}(s) \hookrightarrow GL_{m_0}(q^{2k})$. If $u \in \widetilde{W}_i$ and $v \in \widetilde{W}_j$ then $\tau(u,v) = (su, sv) = \lambda^{q^{i-1}+q^{j-1}}(u,v)$. Therefore,

$$\widetilde{W}_1^{\perp} = \bigoplus_{i \neq 1+k} \widetilde{W}_i \text{ and } \widetilde{W}_{1+k} \cap \widetilde{W}_1^{\perp} = 0. \tag{2.2}$$

Choose a basis $(u_1, ..., u_{m_0})$ in $\widetilde{W}_1$. Then (v_i) is a basis for $\widetilde{W}_{1+k}$, where $v_i = \sigma^k(u_i)$. We see that $(\sigma u_i, \sigma v_j) = (u_i, v_j)^q$. Hence, $(u_i, v_j)^{q^k} = (\sigma^k u_i, \sigma^k v_j) = (v_i, u_j) = (u_j, v_i)$. In other words, ${}^tU = U^{q^k}$. Thus, together with 2.2, U determines a non-degenerate Hermitian form on an m_0-dimensional $\mathbb{F}_{q^{2k}}$-space.

If g acts on $\widetilde{W}_1$ with matrix $A = (a_{ij})$ (with respect to the basis (u_i)), then g acts on $\widetilde{W}_{1+k}$ with matrix $A^{q^k} = (a_{ij}^{q^k})$ (with respect to the basis (v_i)). From 2.2, $g \in C_{O(W)}(s)$ if and only if

$$(u_k, v_l) = (gu_k, gv_l) = (\sum_i a_{ik}u_i, \sum_j a_{jl}^{q^k} v_j) = \sum_{i,j} a_{ik}a_{jl}^{q^k}(u_i, v_j),$$

i.e. ${}^tAUA^{q^k} = U$. Therefore, $C_{O(W)}(s) \simeq GU_{m_0}(q^k)$. Similarly, $g \in C_{CO(W)}(s)$ if and only if ${}^tAUA^{q^k} = \tau(g)U$. Note that if ${}^tAUA^{q^k} = U$ then ${}^t(\varepsilon A)U(\varepsilon A)^{q^k} = \tau(g)U$, where ε is a scalar in $\mathbb{F}_{q^{2k}}$ such that $\varepsilon^{q^k+1} = \tau(g)$ (there always exists such an ε for any $\tau(g) \in \mathbb{F}_q^*$). That means $\tau : C_{CO(W)}(s) \longrightarrow \mathbb{F}_q^*$ is an epimorphism and hence $C_{CO(W)}(s) \simeq C_{O(W)}(s) \cdot \mathbb{Z}_{q-1}$.

1b) Next, we show that $C_{O(W)}(s) \simeq GL_{n_0}(q^k)$ and $C_{CO(W)}(s) \simeq GL_{n_0}(q^k) \cdot \mathbb{Z}_{q-1}$, where $W = U_j$ for $j = 1, ..., l'$, $n_0 = n_j$, and $k = \deg(g_j)$. Note that the characteristic polynomial of s acting on W is $g_j^{n_0}(x)\widehat{g}_j^{n_0}(x)$. Let $\lambda \in \mathbb{F}_{q^k}$ be a root of g_j. Then all the roots of g_j are $\lambda, \lambda^q, ..., \lambda^{q^{k-1}}$ and all the roots of $\widehat{g}_j$ are $\tau\lambda^{-1}, ..., \tau\lambda^{-q^{k-1}}$. Let (f_i) be a basis of W, and define a Frobenius endomorphism $\sigma : \Sigma_i x_i f_i \mapsto \Sigma_i x_i^q f_i$ on $\widetilde{W} := W \otimes_{\mathbb{F}_q} \mathbb{F}_{q^k}$, where $x_i \in \mathbb{F}_{q^k}$. The simplicity of s implies that $\widetilde{W} = \widetilde{W}_1 \oplus \cdots \oplus \widetilde{W}_k \oplus \widetilde{W}'_1 \oplus \cdots \oplus \widetilde{W}'_k$, where $\widetilde{W}_i = \mathrm{Ker}(s - \lambda^{q^{i-1}})$ and $\widetilde{W}'_j = \mathrm{Ker}(s - \tau\lambda^{-q^{j-1}})$. We see that σ permutes the $\widetilde{W}_i$'s and $\widetilde{W}'_j$'s cyclically: $\sigma(\widetilde{W}_i) = \widetilde{W}_{i+1}$, $\sigma(\widetilde{W}'_j) = \widetilde{W}'_{j+1}$, where $\widetilde{W}_{2k+1} = \widetilde{W}_1$ and $\widetilde{W}'_{2k+1} = \widetilde{W}'_1$. Let $g \in CO(W)$ commuting with $s|_W$. Then g preserves each $\widetilde{W}_i$, $\widetilde{W}'_j$. Again, the action of g on $\widetilde{W}$ is completely determined by its action on $\widetilde{W}_1$ and $\widetilde{W}'_1$. So $C_{CO(W)}(s) \hookrightarrow GL_{2n_0}(q^k)$. We also have

$$\widetilde{W}_1^{\perp} = \bigoplus_i \widetilde{W}_i \bigoplus_{j \geq 2} \widetilde{W}'_j \text{ and } \widetilde{W}'_1 \cap \widetilde{W}_1^{\perp} = 0. \tag{2.3}$$

Choose basis (u_i) and (v_i) of $\widetilde{W}_1$ and $\widetilde{W}'_1$, respectively. Suppose that g acts on $\widetilde{W}_1$ with matrix $A = (a_{ij})$ (with respect to the basis (u_i)) and on $\widetilde{W}'_1$ with matrix (b_{ij})

(with respect to the basis (v_i)). From 2.3, $g \in C_{O(W)}(s)$ if and only if $(gu, gv) = (u, v)$ for any $u \in \widetilde{W}_1$, $v \in \widetilde{W}'_1$, i.e.

$$(u_k, v_l) = (gu_k, gv_l) = (\sum_i a_{ik}u_i, \sum_j b_{jl}v_j) = \sum_{i,j} a_{ik}b_{jl}(u_i, v_j).$$

In other words, $g \in C_{O(W)}(s)$ if and only if ${}^tAUB = U$. Since U is non-degenerate, ${}^tAUB = U$ means that B is uniquely determined by A. Therefore, $C_{O(W)}(s) \simeq GL_{n_0}(q^k)$. Similarly, $g \in C_{CO(W)}(s)$ if and only if $(gu, gv) = \tau(g)(u, v)$ for any $u \in \widetilde{W}_1$, $v \in \widetilde{W}'_1$ or equivalently ${}^tAUB = \tau(g)U$. Therefore, $C_{CO(W)}(s) \simeq C_{O(W)}(s) \cdot \mathbb{Z}_{q-1}$.

1c) Lastly, we show that $C_{O(V_\tau)}(s) \simeq O^{\pm}_m(q^2)$ and $C_{CO(V_\tau)}(s) \simeq C_{O(V_\tau)}(s) \cdot \mathbb{Z}_{q-1}$. Again, we have $\widetilde{V}_\tau = \widetilde{V}_1 \oplus \widetilde{V}_2$, where $\widetilde{V}_\tau = V_\tau \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}$, $\widetilde{V}_1 = \mathrm{Ker}(s - \lambda_1)$, and $\widetilde{V}_2 = \mathrm{Ker}(s + \lambda_1)$. Furthermore, $\widetilde{V}_1 \perp \widetilde{V}_2$ and therefore $\widetilde{V}_1$, $\widetilde{V}_2$ are non-degenerate. Similar arguments as in $1a)$, we see that the action of an element $g \in C_{CO(V_\tau)}(s)$ on $\widetilde{V}_\tau$ is completely determined by its action on $\widetilde{V}_1$. Hence, $C_{O(V_\tau)}(s) \simeq O(\widetilde{V}_1) = O^{\pm}_m(q^2)$ and $C_{CO(V_\tau)}(s) \simeq C_{O(V_\tau)}(s) \cdot \mathbb{Z}_{q-1}$, since m is even.

Combining what we have proved in $1a), 1b), 1c)$, we have that

$$C_{O(V)}(s) \simeq C_{O(V_\tau)}(s) \times \prod_i C_{O(V_i)}(s) \times \prod_j C_{O(U_j)}(s) \simeq$$

$$\simeq O^{\pm}_m(q^2) \times \prod_i GU_{m_i}(q^{\deg(f_i)/2}) \prod_j GL_{n_j}(q^{\deg(g_j)}).$$

We have also proved that $C_{CO(W)}(s) \simeq C_{O(W)}(s) \cdot \mathbb{Z}_{q-1}$ for any $W \in \{V_\tau, V_1, ..., V_l, U_1, ..., U_{l'}\}$. More precisely, given any $\alpha \in \mathbb{F}_q^*$, there exists $g_W \in C_{CO(W)}(s)$ such that $\tau(g_W) = \alpha$ for $W \in \{V_\tau, V_1, ..., V_l, U_1, ..., U_{l'}\}$. Set $g = (g_{V_\tau}, g_{V_1}, ..., g_{V_l}, g_{U_1}, ..., g_{U_{l'}})$. Then $g \in C_{CO(V)}(s)$ and $\tau(g) = \alpha$. Hence, $C_{CO(V)}(s) \simeq C_{O(V)}(s) \cdot \mathbb{Z}_{q-1}$. The Lemma is proved in this case.

Case 2: If τ is a square in $\mathbb{F}_q$, suppose that $\tau = \lambda_2^2$ with $\lambda_2 \in \mathbb{F}_q$. Replacing s by $\lambda_2^{-1}s$ if necessary, we can assume that $\tau = 1$. Decompose $P(x)$ into irreducible polynomials over $\mathbb{F}_q$:

$$P(x) = (x-1)^{k'}(x+1)^{2m-k'} \prod_{i=1}^{l} f_i^{m_i}(x) \prod_{j=1}^{l'} g_j^{n_j}(x)\widehat{g_j}^{n_j}(x),$$

where

- If λ is a root of $f_i(x)$ then λ^{-1} is also a root of $f_i(x)$, $\deg(f_i)$ is even, ± 1 are not the roots of f_i,
- If λ is a root of $g_j(x)$ then λ^{-1} is a root of $\widehat{g_j}(x)$, $\deg(g_j) = \deg(\widehat{g_j})$, $g_j \neq \widehat{g_j}$,
- $n = m + \Sigma_{i=1}^{l} m_i \deg(f_i)/2 + \Sigma_{j=1}^{l'} n_j \deg(g_j)$.

Since $1 = \tau^n = \det(s) = (-1)^{2m-k'}$, k' is even. Set $k' = 2k$ and

$$V' := \mathrm{Ker}(s-1), V'' := \mathrm{Ker}(s+1), V_i := (\frac{P(x)}{f_i^{m_i}(x)})_{x=s}(V), U_j := (\frac{P(x)}{g_j^{n_j}(x)\widehat{g}_j^{n_j}(x)})_{x=s}(V).$$

Then $V = V' \oplus V" \oplus V_1 \oplus \cdots V_l \oplus U_1 \oplus \cdots \oplus U_{l'}$ is an orthogonal decomposition of V. It is obvious that $C_{O(V')}(s) = O(V') = O^{\pm}_{2k}(q)$ and $C_{O(V")}(s) = O(V") = O^{\pm}_{2(m-k)}(q)$. Furthermore, $C_{CO(V')}(s) = CO(V') = O^{\pm}_{2k}(q) \cdot \mathbb{Z}_{q-1}$ and $C_{CO(V")}(s) = CO(V") = O^{\pm}_{2(m-k)}(q) \cdot \mathbb{Z}_{q-1}$. Now we can repeat the arguments in Case 1 and complete the proof. $\square$

Remark. Lemma 2.5 is wrong when $s \notin CO^{\pm}_{2n}(q)^0$. Here is one example. Suppose that q is odd. Let $V = \mathbb{F}_q^2 = \{(x,y) \mid x, y \in \mathbb{F}_q\}$ be the vector space of dimension 2. The quadratic form $Q(x,y) = xy$ is non-degenerate of Witt index 1. The Gram matrix of Q corresponding to the basis $\{(1,0),(0,1)\}$ is $J = \left(\begin{smallmatrix}0 & 1\\ 1 & 0\end{smallmatrix}\right)$. By definition, $O^+_2(q)$ is the group of matrices $A \in GL_2(q)$ such that ${}^tAJA = J$. Therefore,

$$O^+_2(q) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}, \begin{pmatrix} 0 & a \\ a^{-1} & 0 \end{pmatrix} \mid a \in \mathbb{F}_q^* \right\}.$$

Similarly,

$$CO^+_2(q) = \{A \in GL_2(q) \mid \exists \tau \in \mathbb{F}_q^*, {}^tAJA = \tau J\} = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \begin{pmatrix} 0 & a \\ b & 0 \end{pmatrix} \mid a, b \in \mathbb{F}_q^* \right\}.$$

Take the element $s = J$, which is semi-simple. Since $\tau(s) = 1$ and $\det(s) = -1$, $s \notin CO^+_2(q)^0$. Direct computation shows that

$$C_{O^+_2(q)}(s) = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \right\}.$$

On the other hand,

$$C_{CO^+_2(q)}(s) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}, \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix} \mid a \in \mathbb{F}_q^* \right\}.$$

Therefore, $C_{CO^+_2(q)}(s) \ncong C_{O^+_2(q)}(s) \cdot \mathbb{Z}_{q-1}$ by order comparison.

3. UNIPOTENT CHARACTERS

3.1. **Unipotent Characters of $SO_{2n+1}(q)$ and $PCSp_{2n}(q)$.** Proposition 5.1 of [TZ1] shows that $SO_{2n+1}(q)$ as well as $PCSp_{2n}(q)$ have a unique unipotent character of minimal degree $(q^n-1)(q^n-q)/2(q+1)$ and any other non-trivial unipotent character has degree greater than $q^{2n}/2(q+1)$. We mimic its proof and get Proposition 3.1, which classifies unipotent characters of degrees up to $\approx q^{6n-15}/2$.

From [C1, p. 466, 467] , we know that the unipotent characters of $SO_{2n+1}(q)$ and $PCSp_{2n}(q)$ are labeled by symbols of the form

$$\binom{\lambda}{\mu} = \binom{\lambda_1\ \lambda_2\ \lambda_3\ \ldots\ \lambda_a}{\mu_1\ \mu_2\ \ldots\ \mu_b}$$

where $0 \le \lambda_1 < \lambda_2 < \ldots < \lambda_a$, $0 \le \mu_1 < \mu_2 < \ldots < \mu_b$, $a - b$ is odd and positive, $(\lambda_1, \mu_1) \neq (0,0)$, and

$$\sum_i \lambda_i + \sum_j \mu_j - \left(\frac{a+b-1}{2}\right)^2 = n.$$

The integer n is called the rank of the symbol $\binom{\lambda}{\mu}$. The degree of the unipotent character $\chi^{\lambda,\mu}$ corresponding to the symbol $\binom{\lambda}{\mu}$ is equal to

$$\frac{(q^2-1)(q^4-1)\cdots(q^{2n}-1)\prod_{i'<i}(q^{\lambda_i}-q^{\lambda_{i'}})\prod_{j'<j}(q^{\mu_j}-q^{\mu_{j'}})\prod_{i,j}(q^{\lambda_i}+q^{\mu_j})}{2^{\frac{a+b-1}{2}}q^{\binom{a+b-2}{2}+\binom{a+b-4}{2}+\cdots}\prod_i\prod_{k=1}^{\lambda_i}(q^{2k}-1)\prod_j\prod_{k=1}^{\mu_j}(q^{2k}-1)}.$$

Proposition 3.1. *Let G^* be either $(B_n)_{ad}(q) = SO_{2n+1}(q)$ or $(C_n)_{ad}(q) = PCSp_{2n}(q)$. Suppose that $n \ge 6$ and $\chi \in \mathrm{Irr}(G^*)$ is unipotent. Then either χ is one of characters labeled by $\binom{n}{-}$, $\binom{0\ 1\ n}{-}$, $\binom{0\ 1}{n}$, $\binom{1\ n}{0}$, $\binom{0\ n}{1}$, $\binom{0\ 2\ n-1}{-}$, $\binom{0\ 2}{n-1}$, $\binom{2\ n-1}{0}$, $\binom{0\ n-1}{2}$, $\binom{1\ n-1}{1}$, $\binom{0\ 1\ 2\ n}{1}$, $\binom{0\ 1\ 2}{1\ n}$, $\binom{1\ 2\ n}{0\ 1}$, $\binom{0\ 1\ n}{1\ 2}$, or $\chi(1) \ge (q^{2n-2}-1)(q^{2n}-1)(q^{n-2}-1)(q^{n-2}-q^3)/2(q^2-1)(q^4-1)(q^3+1)$ ($\approx q^{6n-15}/2$), which is the degree of the unipotent character labeled by $\binom{0\ 3\ n-2}{-}$.*

Note: For reader's convenience, we put these low-dimensional unipotent characters in Table 1, where the degree of each character is calculated. To keep the paper not too lengthy, we will skip detailed arguments of some inequalities in this and ongoing proofs. We frequently use the following obvious estimates without notice:

(i) $q^{a-b} \le \dfrac{q^a-1}{q^b-1} \le q^{a-b}(1+\dfrac{1}{q^b-1})$ for $a \ge b \ge 1$.

(ii) $\prod_{i=2}^n \dfrac{q^{a_i}-q^{a_1}}{q^{a_i}-q^{a_1-1}} > \dfrac{1}{2}$ for integers $1 \le a_1 < a_2 < \ldots < a_n$.

Proof. 1) Define

$$D(n) = (q^{2n-2}-1)(q^{2n}-1)(q^{n-2}-1)(q^{n-2}-q^3)/2(q^2-1)(q^4-1)(q^3+1).$$

We have that

$$\frac{D(n)}{D(n-1)} = \frac{(q^{2n}-1)(q^{n-2}-1)(q^{n-5}-1)}{(q^{2n-4}-1)(q^{n-3}-1)(q^{n-6}-1)}$$

and $q^6 < D(n)/D(n-1) < q^7$ for every $n \ge 7$. Also, $D(n) < q^{6n-14}/2$ for every $n \ge 6$. Let $\mathcal{L}_n$ be the set of 14 symbols in this Proposition. Note that if $(n,q) \neq (6,2)$ then the degrees of characters corresponding to the symbols in $\mathcal{L}_n$ is smaller than $D(n)$. We will prove by induction on $n \ge 6$ that if $\chi = \chi^{\lambda,\mu}$ with $\binom{\lambda}{\mu}$ is not in $\mathcal{L}_n$, then $\chi(1) \geq D(n)$. The induction base $n = 6$ can be checked easily by using [CH, Table 27]. The rest of the proof establishes the induction step for $n \geq 7$, by means of induction on $b \geq 0$.

2) Now we consider the case $b = 0$. Then $a \geq 1$ is odd. We may assume $a \geq 3$, and $\lambda \neq (0, 1, n)$, $(0, 2, n-1)$, $(0, 3, n-2)$. First we assume $a = 3$. If $\lambda_1 = 0$ then $\lambda = (0, k, n+1-k)$, where $4 \leq k < (n+1)/2$; in particular, $n \geq 8$. In this case,

$$\chi(1) = \frac{1}{2} \cdot \frac{(q^{2(k+1)} - 1) \cdots (q^{2n} - 1)}{(q^4 - 1) \cdots (q^{2(n-k+1)} - 1)} \cdot \frac{(q^k - 1)(q^{n-k+1} - 1)(q^{n-k+1} - q^k)}{q^2 - 1} >$$

$$> \frac{1}{2} q^{2(k-1)(n-k)} \cdot \frac{(q^k - 1)(q^{n-k+1} - 1)(q^{n-k+1} - q^k)}{q^2 - 1}.$$

Since $3 \leq k - 1 < n - k$, $2(k-1)(n-k) \geq 6(n-4) = 6n - 24$. Moreover, $n - k + 1 > k$. Therefore $(q^k - 1)(q^{n-k+1} - 1)(q^{n-k+1} - q^k)/(q^2 - 1) > q^{k-2} q^{2k} \geq q^{10}$. It follows that $\chi(1) > \frac{1}{2} q^{6n-24} q^{10} = q^{6n-14}/2 > D(n)$ as required. If $\lambda_1 \geq 1$, then $\lambda = (k, l, n+1-k-l)$ with $1 \leq k < l < n+1-k-l$. Then

$$\chi(1) = \frac{1}{2} \cdot \frac{(q^{2(k+1)} - 1) \cdots (q^{2(k+l)} - 1)}{(q^2 - 1) \cdots (q^{2l} - 1)} \cdot \frac{(q^{2(k+l+1)} - 1) \cdots (q^{2n} - 1)}{(q^4 - 1) \cdots (q^{2(n+1-k-l)} - 1)} \cdot$$

$$\cdot \frac{(q^l - q^k)(q^{n+1-k-l} - q^k)(q^{n+1-k-l} - q^l)}{q^2 - 1} > \frac{1}{2} q^{2kl} q^{2(k+l-1)(n-k-l)} q^{2k+l}.$$

If $(k, l) = (1, 2)$ then $\lambda = (1, 2, n-2)$ and

$$\chi(1) = \frac{(q^2 - q)(q^{2n-2} - 1)(q^{2n} - 1)(q^{n-2} - q)(q^{n-2} - q^2)}{2(q^2 - 1)^2 (q^4 - 1)} > D(n)$$

for every $n \geq 6$. It remains to consider $(k, l) \neq (1, 2)$. Then $k + l \geq 4$ and $l \geq 3$. It follows that $3 \leq k + l - 1$ and $3 \leq l \leq n - k - l$. Therefore, $2(k + l - 1)(n - k - l) \geq 6(n-4)$. So, in this case, $\chi(1) > \frac{1}{2} q^{2kl} q^{6n-24} q^{2k+l} > \frac{1}{2} q^{6n-14} > D(n)$ and we are done. Now we may assume $a \geq 5$.

Suppose that $\lambda_1 \geq 1$. Consider the unipotent character χ' labeled by the symbol $\binom{\lambda'}{-}$ of rank $n - 1$, where $\lambda' = (\lambda_1 - 1, \lambda_2, ..., \lambda_a)$. We have

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^{\lambda_2} - q^{\lambda_1}) \cdots (q^{\lambda_a} - q^{\lambda_1})}{(q^{\lambda_2} - q^{\lambda_1 - 1}) \cdots (q^{\lambda_a} - q^{\lambda_1 - 1})} \cdot \frac{q^{2n} - 1}{q^{2\lambda_1} - 1} \geq \frac{q^{2(n-\lambda_1)}}{2}.$$

Note that $\sum_{i=1}^{a} \lambda_i - \left(\frac{a-1}{2}\right)^2 = n$ and hence $n - \lambda_1 = \lambda_2 + \cdots + \lambda_a - \left(\frac{a-1}{2}\right)^2 \geq (a-1)\lambda_1 + \frac{a^2-1}{4} \geq 10$. Therefore, $\chi(1)/\chi'(1) \geq q^{20}/2 > q^7$. It follows that $\chi(1) \geq q^7 D(n-1) > D(n)$. Now we may assume that $\lambda_1 = 0$.

Next we assume that $\lambda_i - \lambda_{i-1} \geq 2$ for some $i \geq 2$. Consider the unipotent character χ' labeled by the symbol $\binom{\lambda'}{-}$ of rank $n - 1$, where $\lambda' = (\lambda_1, ..., \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, ..., \lambda_a)$. Similarly, we have $\chi(1)/\chi'(1) > q^{2(n-\lambda_i)}/2$. By induction hypotheses, $\chi'(1) \geq D(n-1)$. If $n - \lambda_i \geq 4$ then $\chi(1) \geq q^7 D(n-1) > D(n)$. Now suppose

$n-\lambda_i \le 3$. Note that $\sum_i \lambda_i = \left(\frac{a-1}{2}\right)^2 + n$. If $i \le a-1$ then $\lambda_{a-1} \ge n-3$ and $\lambda_a \ge n-2$. Then $\left(\frac{a-1}{2}\right)^2 + n \ge 0+1+\cdots+(a-3)+(n-3)+(n-2)$ or $a^2-8a-9+4n \le 0$. This is a contradiction because $a \ge 5$ and $n \ge 7$. Therefore $i=a$ and we have $0+1+\cdots+(a-2)+\lambda_a = \left(\frac{a-1}{2}\right)^2 + n$. Since $\lambda_a \ge (n-3)$, $a^2-4a-9 \le 0$ and therefore $a=5$. Then $\lambda = (0,1,2,3,n-2)$ and it is easy to check that $\chi(1) > D(n)$. Now we may assume $\lambda = (0,1,2,...,a-2,a-1)$. Consider $\chi' = \chi^{\lambda',\mu'}$, where $\lambda' = (0,1,2,...,a-4,a-3)$, and $\mu' = \mu$. Note that $a^2 = 4n+1$ and the rank of χ' is $n-a+1$. We have

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^{2(n-a+2)}-1)...(q^{2n}-1)\prod_{i\ge a-2, i>i'}(q^i-q^{i'})}{2q^{\binom{a-2}{2}}\prod_{k=1}^{a-2}(q^{2k}-1)\prod_{k=1}^{a-1}(q^{2k}-1)} =$$

$$= \frac{(q^{2(n-a+2)}-1)...(q^{2n}-1)}{(q^2-1)...(q^{2(a-1)}-1)} \cdot \frac{q^{a-1}-q^{a-2}}{2} \cdot \prod_{i=0}^{a-3} \frac{(q^{a-1}-q^i)(q^{a-2}-q^i)}{q^i(q^{2(i+1)}-1)}.$$

Remark that

$$\prod_{i=0}^{a-3} \frac{(q^{a-1}-q^i)(q^{a-2}-q^i)}{q^i(q^{2(i+1)}-1)} = \prod_{i=0}^{a-3} \frac{q^i(q^{i+2}-1)}{q^{i+1}+1} \ge \prod_{i=0}^{2} \frac{q^i(q^{i+2}-1)}{q^{i+1}+1} > q^4.$$

Therefore,

$$\frac{\chi(1)}{\chi'(1)} \ge \frac{q^{2(n-a+1)(a-1)}(q^{a-1}-q^{a-2})q^4}{2}.$$

Since $n \ge 7$ and $a^2 = 4n+1$, $n \ge 12$ and $a \ge 7$. Then $(n-a+1)(a-1) \ge 6(n-6)$. Hence, $\chi(1)/\chi'(1) > q^{12n-63}/2 > q^7$ and we are done.

3) From now on we assume that $b \ge 1$. At this point we suppose that $(\lambda_1,\mu_1) \ne (1,0)$ and $\lambda_1 \ge 1$. Consider the character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda' = (\lambda_1-1,\lambda_2,...,\lambda_a)$ and $\mu' = \mu$. If $\binom{\lambda'}{\mu'} \in \mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu} \in \left\{\binom{1\ 2}{n-2}, \binom{3\ n-2}{0}, \binom{1\ n-2}{2}, \binom{2\ n-2}{1}\right\}$. It is easy to check that the degrees of characters corresponding to these symbols are greater than $D(n)$. Now we can assume $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. We have

$$\frac{\chi(1)}{\chi'(1)} = \frac{q^{2n}-1}{q^{2\lambda_1}-1} \cdot \prod_{i=2}^{a} \frac{q^{\lambda_i}-q^{\lambda_1}}{q^{\lambda_i}-q^{\lambda_1-1}} \cdot \prod_{j=1}^{b} \frac{q^{\mu_j}+q^{\lambda_1}}{q^{\mu_j}+q^{\lambda_1-1}} > \frac{q^{2n}-1}{2(q^{2\lambda_1}-1)} > \frac{q^{2(n-\lambda_1)}}{2}.$$

If $n-\lambda_1 \le 3$ then $\lambda_1 \ge n-3$ and we have $n-3+n-2+...+n+a-4+\frac{(b-1)b}{2} \le \frac{(a+b-1)^2}{4}+n$. This implies $4a(n-3)+(a-b)^2 \le 4n+1$ and therefore $a(n-3) \le n$. Since $n \ge 7$, $a=1$ which leads to a contradiction since $b \ge 1$. So $n-\lambda_1 \ge 4$. Then we have $\chi(1)/\chi'(1) > q^7$ and $\chi(1) > q^7D(n-1) > D(n)$ as desired.

Similarly, for $(\lambda_1, \mu_1) \neq (0, 1)$ and $\mu_1 \geq 1$, we consider the character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda' = \lambda$ and $\mu' = (\mu_1 - 1, \mu_2, ..., \mu_b)$. If $\binom{\lambda'}{\mu'} \in \mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu} \in \left\{\binom{2\ n-2}{1}, \binom{0\ n-2}{3}, \binom{1\ n-2}{2}, \binom{0\ 1\ 2\ n-1}{2}, \binom{0\ 1\ 2}{2\ n-1}\right\}$. Again, it is easy to check that the degrees of characters corresponding to these symbols are greater than $D(n)$. Now we can assume $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. We have

$$\frac{\chi(1)}{\chi'(1)} = \frac{q^{2n}-1}{q^{2\mu_1}-1} \cdot \prod_{j=2}^{b} \frac{q^{\mu_j} - q^{\mu_1}}{q^{\mu_j} - q^{\mu_1 - 1}} \cdot \prod_{i=1}^{a} \frac{q^{\lambda_i} + q^{\mu_1}}{q^{\lambda_i} + q^{\mu_1 - 1}} > \frac{q^{2n}-1}{2(q^{2\mu_1}-1)} > \frac{q^{2(n-\mu_1)}}{2}.$$

If $n - \mu_1 \geq 4$ then $\chi(1)/\chi'(1) > q^7$ and $\chi(1) > q^7 D(n-1) > D(n)$ as desired. If $n - \mu_1 \leq 3$ then $\mu_1 \geq n-3$ and we have $\frac{(a-1)a}{2} + n - 3 + n - 2 + ... + n + b - 4 \leq \frac{(a+b-1)^2}{4} + n$. This implies $4b(n-3) + (a-b)^2 \leq 4n + 1$. So $b(n-3) \leq n$. Since $n \geq 7$, $b = 1$. Then we have $(a-1)^2 \leq 13$ and therefore $a = 2$ or $a = 4$.

$\mathbf{Case\ a = 4, b = 1}$: Then $\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 + \mu_1 = n + 4$. This implies $\mu_1 \leq n - 2$. Hence $\binom{\lambda}{\mu} = \binom{0\ 1\ 2\ 3}{n-2}$ or $\binom{0\ 1\ 2\ 4}{n-3}$. The degrees of characters corresponding to both these symbols are greater than $D(n)$.

$\mathbf{Case\ a = 2, b = 1}$: Since $n - 3 \leq \mu_1 \leq n$ and $\binom{\lambda}{\mu}$ is not in $\mathcal{L}_n$, we have $\binom{\lambda}{\mu} = \binom{0\ 4}{n-3}$, $\binom{1\ 3}{n-3}$, $\binom{0\ 3}{n-2}$ or $\binom{1\ 2}{n-2}$. Again, the degrees of characters corresponding to these symbols are greater than $D(n)$.

It remains to consider the case where $(\lambda_1, \mu_1) = (0, 1)$ or $(1, 0)$.

4) Here we suppose that $(\lambda_1, \mu_1) = (1, 0)$ but $\lambda \neq (1, 2, ..., a)$. Then there exists an $i \geq 2$ such that $\lambda_i \geq \lambda_{i-1} + 2$. We choose i to be smallest possible. If $a = 2$ then $b = 1$ and $\binom{\lambda}{\mu} = \binom{1\ n}{0} \in \mathcal{L}_n$. Hence $a \geq 3$. Consider the character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda' = (\lambda_1, ..., \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, ..., \lambda_a)$ and $\mu' = \mu$. If $\binom{\lambda'}{\mu'} \in \mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu}$ can only be $\binom{1\ 3\ n-1}{0\ 1}$. The degree of the corresponding character is greater than $D(n)$. Now we can assume that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. By induction hypothesis, $\chi'(1) \geq D_{n-1}$. Set

$$T_1 = \prod_{i'<i} \frac{q^{\lambda_i} - q^{\lambda_{i'}}}{q^{\lambda_i - 1} - q^{\lambda_{i'}}}, T_2 = \prod_{i'>i} \frac{q^{\lambda_{i'}} - q^{\lambda_i}}{q^{\lambda_{i'}} - q^{\lambda_i - 1}}, T_3 = \prod_{j} \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i - 1} + q^{\mu_j}}.$$

Then $\chi(1)/\chi'(1) = [(q^{2n}-1)/(q^{2\lambda_i}-1)]T_1 T_2 T_3 > q^{2(n-\lambda_i+1)}/2$ (see [TZ1, p. 2121]). If $n - \lambda_i \geq 3$ then $\chi(1)/\chi'(1) > q^7$ and therefore $\chi(1) \geq D(n)$ as desired. Now we assume $n - \lambda_i \leq 2$. There are two cases:

$\mathbf{Case\ n - \lambda_i = 2}$: Then $i \geq a - 1$. If $i = a - 1$ then $a - b = 1$, $a = n - 2$ and $\binom{\lambda}{\mu} = \binom{1\ \cdots\ n-4\ n-2\ n-1}{0\ \cdots\ n-4}$. Since $n \geq 7$, $a \geq 5$ and $i = b \geq 4$. It follows that $T_1 \geq q^3$, $T_3 \geq q^{b-1} \geq q^3$. Hence $\chi(1)/\chi'(1) > q^{10}/2 > q^7$ and we are done. If $i = a \geq 3$ then $T_1 \geq q^{a-2}\frac{q^{\lambda_i}-q}{q^{\lambda_i - 1}-q}$, $T_2 = 1$ and $T_3 \geq \frac{q^{\lambda_a}+1}{q^{\lambda_a - 1}+1}$. Then $\chi(1)/\chi'(1) > q^7$ and we are done.

Case $\mathbf{n}-\lambda_{\mathbf{i}} \leq \mathbf{1}$: Then $i=a$ and $\lambda_a \geq n-1$. If $\lambda_a \geq n$ then $\lambda_a = n$, $a-b=1$ and $\binom{\lambda}{\mu} = \binom{1\ 2\ \dots\ a-1\ n}{0\ 1\ \dots\ a-2}$. Since $\binom{\lambda}{\mu}$ is not in $\mathcal{L}_n$, $a \geq 4$. We have

$$\frac{\chi(1)}{\chi'(1)} = \prod_{j=1}^{a-1} \left(\frac{q^n - q^j}{q^{n-1} - q^j} \cdot \frac{q^n + q^{j-1}}{q^{n-1} + q^{j-1}} \right) = \frac{(q^n+1)(q^{n-1}-1)}{(q^{n-a}-1)(q^{n-a+1}+1)} > q^{2a-2}.$$

If $a \geq 5$ then $\chi(1)/\chi'(1) > q^8$ and therefore we are done. If $a=4$ then $\binom{\lambda}{\mu} = \binom{1\ 2\ 3\ n}{0\ 1\ 2}$ and $\chi(1) > D(n)$. If $\lambda_a = n-1$ then we also have $a-b=1$ and $\binom{\lambda}{\mu} = \binom{1\ 2\ \dots\ a-1\ n-1}{0\ 1\ \dots\ a-3\ a-1}$. If $a=3$ then $\binom{\lambda}{\mu} = \binom{1\ 2\ n-1}{0\ 2}$. Check directly we see that $\chi(1) > D(n)$. If $a \geq 4$ then $T_1 > q^{a-1} \geq q^3$, $T_2 = 1$ and $T_3 \geq q^{b-1} \geq q^2$. So $\chi(1)/\chi'(1) > q^7$ and we are done again.

Similarly, suppose that $(\lambda_1, \mu_1) = (0,1)$ but $\mu \neq (1,2,...,b)$; in particular, $b \geq 2$. Then there exists an index $j \geq 2$ such that $\mu_j \geq \mu_{j-1}+2$. We choose j to be smallest possible. Consider the character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda' = \lambda$ and $\mu' = (\mu_1, ..., \mu_{j-1}, \mu_j - 1, \mu_{j+1}, ..., \mu_b)$. If $\binom{\lambda'}{\mu'} \in \mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu}$ can only be $\binom{0\ 1\ n-1}{1\ 3}$. The degree of the corresponding character is greater than $D(n)$ and hence we are done in this case. Now we can assume $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. Set

$$U_1 = \prod_{j'<j} \frac{q^{\mu_j} - q^{\mu_{j'}}}{q^{\mu_j - 1} - q^{\mu_{j'}}}, U_2 = \prod_{j'>j} \frac{q^{\mu_{j'}} - q^{\mu_j}}{q^{\mu_{j'}} - q^{\mu_j - 1}}, U_3 = \prod_i \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i} + q^{\mu_j - 1}}.$$

Then $\chi(1)/\chi'(1) = [(q^{2n}-1)/(q^{2\mu_j}-1)]U_1U_2U_3 > q^{2(n-\mu_j+1)}/2$ (see [TZ1, p. 2122]). If $n - \mu_j \geq 3$ then $\chi(1)/\chi'(1) > q^7$ and therefore $\chi(1) \geq D_n$ as desired. Now we assume $n - \mu_j \leq 2$. There are two cases:

Case $\mathbf{n}-\mu_{\mathbf{j}} = \mathbf{2}$: Then $j \geq b-1$. If $j = b-1$ then $a-b=1$, $b=n-2$ and $\binom{\lambda}{\mu} = \binom{0\ 1\ \dots\ n-2}{1\ \dots\ n-4\ n-2\ n-1}$. Since $n \geq 7$, $b \geq 5$ and $j \geq 4$. It follows that $U_1 \geq q^2 \frac{q^{\mu_j}-q}{q^{\mu_j-1}-q}$, $T_3 \geq \frac{q^{\mu_j}+1}{q^{\mu_j-1}+1}$. Hence $\chi(1)/\chi'(1) > q^8/2 > q^7$ and we are done. If $j = b \geq 3$ then $V_1 \geq q\frac{q^{\mu_j}-q}{q^{\mu_j-1}-q}$, $V_2 = 1$ and $V_3 \geq \frac{q^{\lambda_a}+1}{q^{\lambda_a-1}+1}$. Therefore, $\chi(1)/\chi'(1) > q^7$ and we are done again. If $j=b=2$ then $a=3$ or $a=5$. In any case, $\lambda_a \leq 4 < n-2$. Therefore, in this case, $U_1 > q$, $U_2 = 1$ and $U_3 > q^2$. Then $\chi(1)/\chi'(1) > q^7$ and $\chi(1) \geq D_n$ as required.

Case $\mathbf{n}-\mu_{\mathbf{j}} = \mathbf{1}$: Then $j = b$ and $\mu_b \geq n-1$. If $\mu_b \geq n$ then $\mu_b = n$, $a-b=1$ and $\binom{\lambda}{\mu} = \binom{0\ 1\ \dots\ b}{1\ 2\ \dots\ b-1\ n}$. Since $\binom{\lambda}{\mu}$ is not in $\mathcal{L}_n$, $b \geq 3$. We have

$$\frac{\chi(1)}{\chi'(1)} = \prod_{i=0}^{b} \frac{q^n + q^i}{q^{n-1} + q^i} \cdot \prod_{i=1}^{b-1} \frac{q^n - q^i}{q^{n-1} - q^i} = \frac{(q^n+1)(q^{n-1}-1)}{(q^{n-b-1}+1)(q^{n-b}-1)} > q^{2b-1}.$$

If $b \geq 4$ then $\chi(1)/\chi'(1) > q^7$ and therefore we are done. If $b=3$ then $\binom{\lambda}{\mu} = \binom{0\ 1\ 2\ 3}{1\ 2\ n}$. In this case, $\chi(1) > D(n)$. If $\mu_b = n-1$ then we also have $a-b=1$ and $\binom{\lambda}{\mu} =$

$\left(\begin{smallmatrix}0 & 1 & \dots & b-1 & b+1\\ 1 & 2 & \dots & b-1 & n-1\end{smallmatrix}\right)$ where $n \geq b+2$. If $b=2$ then $\binom{\lambda}{\mu} = \left(\begin{smallmatrix}0 & 1 & 3\\ 1 & n-1 & \end{smallmatrix}\right)$. Check directly we see that $\chi(1) > D(n)$. If $b \geq 3$ then $U_1 > q^{b-1} \geq q^2$, $U_2 = 1$ and $U_3 \geq q^3$. So $\chi(1)/\chi'(1) > q^7$ and we are done.

5) Here we suppose that $\mu_1 = 0$ and $\lambda = (1, 2, ..., a)$. Consider the character χ' labeled by the symbol

$$\binom{\lambda'}{\mu'} = \begin{pmatrix} 0 & 1 & 2 & \dots & a \\ \mu_2 & \mu_3 & \dots & \mu_b & \end{pmatrix}$$

of the same rank n, but with the parameters $a' = a+1$ and $b' = b-1$. It is easy to see that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_n$. So $\chi'(1) > D(n)$ by the induction hypothesis (on b). But

$$\frac{\chi(1)}{\chi'(1)} = \frac{\prod_{i=1}^{a}(1+\frac{2}{q^i-1})}{\prod_{j=2}^{b}(1+\frac{2}{q^{\mu_j}-1})} \geq \frac{\prod_{i=1}^{a}(1+\frac{2}{q^i-1})}{\prod_{j=1}^{b-1}(1+\frac{2}{q^j-1})} > 1,$$

so we are done.

Similarly, suppose that $\lambda_1 = 0$ and $\mu = (1, 2, ..., b)$ but $\lambda \neq (0, 1, ..., a-1)$. Then there exists an $i \geq 2$ such that $\lambda_i \geq \lambda_{i-1} + 2$. We choose i to be smallest possible. Consider the character χ' labeled by the symbol

$$\binom{\lambda'}{\mu'} = \begin{pmatrix} \lambda_1 & \dots & \lambda_{i-1} & \lambda_i - 1 & \lambda_{i+1} & \dots & \lambda_a \\ & & & \mu_1 \ \dots \ \mu_b & & & \end{pmatrix}$$

of rank $n' = n-1$. If $\binom{\lambda'}{\mu'} \in \mathcal{L}_{n-1}$, then $\binom{\lambda}{\mu}$ is one of the symbols $\left(\begin{smallmatrix}0 & 1 & 3 & n-1\\ & & 1 & \end{smallmatrix}\right)$, $\left(\begin{smallmatrix}0 & 2 & n-1\\ & 1 & 2\end{smallmatrix}\right)$. In this case, we can check that $\chi(1) > D(n)$. Now we can suppose that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. We have $\frac{\chi(1)}{\chi'(1)} = \frac{q^{2n}-1}{q^{2\lambda_i}-1} \cdot V_1V_2V_3$, where

$$V_1 = \prod_{i'<i} \frac{q^{\lambda_i} - q^{\lambda_{i'}}}{q^{\lambda_i - 1} - q^{\lambda_{i'}}}, V_2 = \prod_{i'>i} \frac{q^{\lambda_{i'}} - q^{\lambda_i}}{q^{\lambda_{i'}} - q^{\lambda_i - 1}}, V_3 = \prod_{j} \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i - 1} + q^{\mu_j}}.$$

Note that $V_1 \geq (q^{\lambda_i} - 1)/(q^{\lambda_i - 1} - 1) > q$, $V_2 > 1/2$ and $V_3 > 1$. So $\chi(1)/\chi'(1) > q^{2(n-\lambda_i)+1}/2$. If $n - \lambda_i \geq 4$, then $\chi(1)/\chi'(1) > q^9/2$ and we are done. Now we assume that $n - \lambda_i \leq 3$. Then $i \geq a-1$. There are two cases:

Case $\mathbf{i = a - 1}$: We have $(a-b)^2 + 4b + 4\lambda_{a-1} + 4\lambda_a = 4n + 8a - 11$. Since $(a-b)^2 + 4b \geq 4a - 3$ and $\lambda_a \geq a$, $\lambda_{a-1} \leq n-2$. If $\lambda_{a-1} = n-2$ then $\lambda_a = a$. It follows that $a \geq n-1 \geq 6$. Hence, $V_1 \geq q^{a-2} \geq q^4$ and we have

$$\frac{\chi(1)}{\chi'(1)} > \frac{1}{2}q^4 \frac{q^{2n}-1}{q^{2n-4}-1} \geq q^7,$$

so $\chi(1) > D_n$. If $\lambda_{a-1} = n-3$ then $\lambda_a \leq a+1$. Then $a+1 \geq n-2$ and therefore $a \geq 4$. Hence $V_1 \geq q^{a-2} \geq q^2$. Again, we have

$$\frac{\chi(1)}{\chi'(1)} > \frac{1}{2}q^2 \frac{q^{2n}-1}{q^{2n-6}-1} \geq q^7$$

and we are done.

Case $\mathbf{i = a}$: Then $(a-b)^2+4b+4\lambda_a = 4n+4a-3$. Therefore, $(a-b-2)^2 = 4(n-\lambda_a)+1$. Since $n-\lambda_a \leq 3$, $n-\lambda_a = 0$ or 2. If $n-\lambda_a = 2$ then $(a-b-2)^2 = 9$ and we have $a-b=5$. In particular, $a \geq 6$. Then $V_1 > q^{a-1} \geq q^5$. Moreover, in this case, $V_2 = 1$ and $V_3 > 1$. So $\chi(1)/\chi'(1) > q^9$ and we are done. If $n-\lambda_a = 0$ then $a-b = 1$ or 3. In this case, $V_1 > q^{a-1}$, $V_2 = 1$ and $V_3 > q^{b-1}$. So $\chi(1)/\chi'(1) > q^{a+b-2}$. If $(a,b) = (4,3)$ or $(5,2)$ then $\binom{\lambda}{\mu} = \binom{0\ 1\ 2\ n}{1\ 2\ 3}$ or $\binom{0\ 1\ 2\ 3\ n}{1\ 2\ 3}$. Checking directly we have $\chi(1) > D(n)$. Otherwise, $a+b \geq 9$ and therefore $\chi(1)/\chi'(1) \geq q^7$ and we are done again.

6) Finally, we consider the case where $\binom{\lambda}{\mu} = \binom{0\ 1\ 2\ \dots\ a-1}{1\ 2\ \dots\ b}$. Consider the character χ' labeled by the symbol $\binom{\lambda'}{\mu'} = \binom{0\ 1\ 2\ \dots\ a-2}{1\ 2\ \dots\ b-1}$ of rank $n-1$. Then we have

$$\frac{\chi(1)}{\chi'(1)} = \frac{q^{a+b-2}(q^{2n}-1)}{(q^b-1)(q^{a-1}+1)} > \frac{q^{2n}-1}{q} > q^7.$$

Note that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. Therefore, $\chi(1) > q^7\chi'(1) \geq q^7 D(n-1) > D(n)$ as desired. ☐

Corollary 3.2. *Let G^* be either $(B_n)_{ad}(q) = SO_{2n+1}(q)$ or $(C_n)_{ad}(q) = PCSp_{2n}(q)$. Suppose that $n \geq 5$ and $\chi \in \mathrm{Irr}(G^*)$ is unipotent. Then either χ is one of characters labeled by $\binom{n}{-}$, $\binom{0\ 1\ n}{-}$, $\binom{0\ 1}{n}$, $\binom{1\ n}{0}$, $\binom{0\ n}{1}$, with degrees 1, $(q^n-1)(q^n-q)/2(q+1)$, $(q^n+1)(q^n+q)/2(q+1)$, $(q^n+1)(q^n-q)/2(q-1)$, $(q^n-1)(q^n+q)/2(q-1)$, respectively, or $\chi(1) > q^{4n-8}$.*

Proof. If $n \geq 6$, Corollary comes from Proposition 3.1. The case $n=5$ can be verified easily by using Table 26 of [CH]. ☐

3.2. **Unipotent Characters of $P(CO_{2n}^-(q)^0)$.**

Proposition 7.1 in [TZ1] shows that the projective conformal orthogonal group of type $-$, $P(CO_{2n}^-(q)^0)$, has a unique unipotent character of minimal degree $(q^n+1)(q^{n-1}-q)/(q^2-1)$ and any other nontrivial unipotent character has degree greater than q^{2n-2}. We mimic its proof and get Proposition 3.3, which classifies unipotent characters of degrees up to q^{4n-10}.

From [C1, p. 475, 476], we know that the unipotent characters of $P(CO_{2n}^-(q)^0)$ are labeled by symbols of the form

$$\binom{\lambda}{\mu} = \binom{\lambda_1\ \lambda_2\ \lambda_3\ \dots\ \lambda_a}{\mu_1\ \mu_2\ \dots\ \mu_b},$$

where $0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_a$, $0 \leq \mu_1 < \mu_2 < \dots < \mu_b$, $a > b$, $a-b \equiv 2 \pmod 4$, $(\lambda_1, \mu_1) \neq (0,0)$, and

$$\sum_i \lambda_i + \sum_j \mu_j - \left[\left(\frac{a+b-1}{2}\right)^2\right] = n.$$

The integer n is called the rank of the symbol $\binom{\lambda}{\mu}$. The degree of the unipotent character $\chi^{\lambda,\mu}$ corresponding to the symbol $\binom{\lambda}{\mu}$ is equal to

$$\frac{(q^n+1)\prod_{i=1}^{n-1}(q^{2i}-1)\prod_{i'<i}(q^{\lambda_i}-q^{\lambda_{i'}})\prod_{j'<j}(q^{\mu_j}-q^{\mu_{j'}})\prod_{i,j}(q^{\lambda_i}+q^{\mu_j})}{2^{\frac{a+b-2}{2}}q^{\binom{a+b-2}{2}+\binom{a+b-4}{2}+\cdots}\prod_i\prod_{k=1}^{\lambda_i}(q^{2k}-1)\prod_j\prod_{k=1}^{\mu_j}(q^{2k}-1)}.$$

Proposition 3.3. *Let $G^*=({}^2D_n)_{ad}(q)=P(CO_{2n}^-(q)^0)$. Suppose that $n\geq 6$ and $\chi\in\mathrm{Irr}(G^*)$ is unipotent. Then either χ is one of the characters labeled by $\binom{0\ n}{-}$, $\binom{1\ n-1}{-}$, $\binom{0\ 1\ n}{1}$ with degrees 1, $(q^n+1)(q^{n-1}-q)/(q^2-1)$, $(q^{2n}-q^2)/(q^2-1)$, respectively, or $\chi(1)>q^{4n-10}$. Furthermore, when $n=5$, G^* has one more character of degree $q^2(q^4+1)(q^5+1)/(q+1)$ (corresponding to the symbol $\binom{2\ 3}{-}$), which is less than q^{4n-10}.*

Proof. Denote $\mathcal{L}_n=\binom{0\ n}{-}$, $\binom{1\ n-1}{-}$, $\binom{0\ 1\ n}{1}$ if $n\geq 6$ and $\mathcal{L}_n=\binom{0\ n}{-}$, $\binom{1\ n-1}{-}$, $\binom{0\ 1\ n}{1}$, $\binom{2\ n-2}{-}$ if $n=5$. We will prove by induction on $n\geq 5$ that if $\chi=\chi^{\lambda,\mu}$ and $\binom{\lambda}{\mu}\notin\mathcal{L}_n$, then $\chi(1)>q^{4n-10}$. The induction base $n=5$ can be checked easily by using [CH, Table 31]. The rest of the proof establishes the induction step for $n\geq 6$.

1) At this point we suppose that $(\lambda_1,\mu_1)\neq(1,0)$ and $\lambda_1\geq 1$ (eventually b may be zero). Consider the unipotent character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda'=(\lambda_1-1,\lambda_2,...,\lambda_a)$ and $\mu'=\mu$. If $\binom{\lambda'}{\mu'}\in\mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu}=\binom{2\ n-2}{-}$ and therefore $\chi(1)=(q^{2n-2}-1)(q^n+1)(q^{n-2}-q^2)/(q^2-1)(q^4-1)>q^{4n-10}$. So we can suppose that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. By induction hypothesis, $\chi'(1)>q^{4(n-1)-10}$. Observe that $n=\sum_i\lambda_i+\sum_j\mu_j-\frac{(a+b)^2-2(a+b)}{4}\geq a\lambda_1+\frac{(a-b)^2}{4}$. Since $a\geq 2$ and $a-b\geq 2$, $2\lambda_1<n$. Note that $n\geq 6$. It follows that $\lambda_1\leq n-4$. We have

$$\frac{\chi(1)}{\chi'(1)}=\frac{(q^n+1)(q^{n-1}-1)}{q^{2\lambda_1}-1}\cdot\prod_{i=2}^{a}\frac{q^{\lambda_i}-q^{\lambda_1}}{q^{\lambda_i}-q^{\lambda_1-1}}\cdot\prod_{j=1}^{b}\frac{q^{\lambda_1}+q^{\mu_j}}{q^{\lambda_1-1}+q^{\mu_j}}>$$

$$>\frac{(q^n+1)(q^{n-1}-1)}{2(q^{2\lambda_1}-1)}\geq\frac{(q^n+1)(q^{n-1}-1)}{2(q^{2n-8}-1)}>q^4,$$

and therefore $\chi(1)>q^{4n-10}$.

2) Similarly, supposing $(\lambda_1,\mu_1)\neq(0,1)$ and $\mu_1\geq 1$, consider the unipotent character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda'=\lambda$ and $\mu'=(\mu_1-1,\mu_2,...,\lambda_b)$. If $\binom{\lambda'}{\mu'}\in\mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu}=\binom{0\ 1\ n-1}{2}$ and therefore $\chi(1)=(q^n+1)(q^{n-1}-1)(q^{n-1}-q)(q^{n-1}+q^2)/2(q^2-1)^2>q^{4n-10}$. So we can suppose that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. By induction hypothesis, $\chi'(1)>q^{4(n-1)-10}$. Set

$$T_1=\prod_{j=2}^{b}\frac{q^{\mu_j}-q^{\mu_1}}{q^{\mu_j}-q^{\mu_1-1}},T_2=\prod_{i=1}^{a}\frac{q^{\lambda_i}+q^{\mu_1}}{q^{\lambda_i}+q^{\mu_1-1}}.$$

Then, if $n - \mu_1 \geq 4$,

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n+1)(q^{n-1}-1)}{q^{2\mu_1}-1} T_1 T_2 > \frac{(q^n+1)(q^{n-1}-1)}{2(q^{2\mu_1}-1)} > q^4.$$

If $n - \mu_1 = 3$ then $b = 1$ since $b\mu_1 < n$ and $n \geq 6$. Then $T_1 = 1$ and we still have $\frac{\chi(1)}{\chi'(1)} > \frac{(q^n+1)(q^{n-1}-1)}{q^{2\mu_1}-1} > q^4$. It follows that $\chi(1) > q^{4n-10}$ when $n - \mu_1 \geq 3$. If $n - \mu_1 \leq 2$, then $\binom{\lambda}{\mu} = \binom{0\ 1\ 2}{n-1}, \binom{0\ 1\ 3}{n-2}$. Checking directly, we again have $\chi^{\lambda,\mu}(1) > q^{4n-10}$.

3) Now we consider the case where $b = 0$ and $\lambda_1 = 0$. If $a = 2$ then we can suppose $\lambda = (k, n-k)$ with $3 \leq k < n-k$ ($k = 2$ is considered already in 1)). Then

$$\chi(1) = \frac{(q^n+1)(q^{n-k}-q^k)\prod_{i=k+1}^{n-1}(q^{2i}-1)}{\prod_{i=1}^{n-k}(q^{2i}-1)} >$$
$$> q^{n+k-2} q^{2(k-1)(n-k-1)} \geq q^{n+1} q^{4n-16} > q^{4n-10}.$$

Now we may assume $a \geq 6$. First we suppose that $\lambda \neq (0, 1, ..., a-1)$. Then there exists $i \geq 2$ such that $\lambda_i - \lambda_{i-1} \geq 2$. Consider the character χ' corresponding to the symbol $\binom{\lambda'}{-}$, where $\lambda' = (..., \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, ...)$. Note that $n - \lambda_i \geq (a-2)^2/4 \geq 4$. Therefore, $\chi(1)/\chi'(1) > (q^n+1)(q^{n-1}-1)/2(q^{2\lambda_i}-1) > q^4$. The induction hypothesis $\chi'(1) > q^{4n-14}$ implies that $\chi(1) > q^{4n-10}$.

Next we suppose that $\lambda = (0, 1, ..., a-1)$. Then $\chi(1) = (q^n+1)f(a)$ and $n = a^2/4$, where

$$f(a) = \frac{(q^2-1)(q^4-1)\cdots(q^{(a^2-4)/2}-1)\prod_{0\leq i'\leq i\leq a-1}(q^i - q^{i'})}{2^{\frac{a-2}{2}} q^{\binom{a-2}{2}+\binom{a-4}{2}+\cdots} \prod_{i=0}^{a-1}\prod_{k=1}^{i}(q^{2k}-1)} \tag{3.4}$$

for any even a. It is easy to show that $f(a)/f(a-2) > q^{a(a-1)(a-3)/2}/2^6 > q^{3a^2/4} = q^{3n}$ for $a \geq 6$. Furthermore, $f(4) > 1$. It follows that $\chi(1) > q^{4n-10}$ and we are done.

It remains to consider the case where $b \geq 1$ and $(\lambda_1, \mu_1) = (0, 1)$ or $(1, 0)$.

4) Here we suppose that $(\lambda_1, \mu_1) = (1, 0)$ but $\lambda \neq (1, 2, ..., a)$. Then there exists $i \geq 2$ such that $\lambda_i \geq \lambda_{i-1} + 2$. Choose i to be smallest possible. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = (\lambda_1, ..., \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, ..., \lambda_a)$ and $\mu' = \mu$. By the induction hypothesis, $\chi'(1) > q^{4n-14}$. Set

$$U_1 = \prod_{i'<i} \frac{q^{\lambda_i} - q^{\lambda_{i'}}}{q^{\lambda_i - 1} - q^{\lambda_{i'}}}, U_2 = \prod_{i'>i} \frac{q^{\lambda_{i'}} - q^{\lambda_i}}{q^{\lambda_{i'}} - q^{\lambda_i - 1}}, U_3 = \prod_j \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i - 1} + q^{\mu_j}}.$$

If $n - \lambda_i \geq 3$ then

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n+1)(q^{n-1}-1)}{q^{2\lambda_i}-1} U_1 U_2 U_3 > \frac{(q^n+1)(q^{n-1}-1)}{q^{2\lambda_i}-1} \cdot \frac{q^{\lambda_i} - q}{q^{\lambda_i - 1} - q} \cdot \frac{1}{2} \cdot \frac{q^{\lambda_i}+1}{q^{\lambda_i - 1}+1} >$$
$$> \frac{(q^n+1)(q^{n-1}-1)q^2}{2(q^{2\lambda_i}-1)} \geq \frac{(q^n+1)(q^{n-1}-1)q^2}{2(q^{2n-6}-1)} > q^4,$$

so $\chi(1) > q^{4n-10}$. If $n - \lambda_i \leq 2$ then $i = a$. Note that $n - \lambda_i \geq 1$. As $b \geq 1$, we have $a \geq 3$. Therefore

$$U_1 \geq \frac{(q^{\lambda_a} - q)(q^{\lambda_a} - q^2)}{(q^{\lambda_a - 1} - q)(q^{\lambda_a - 1} - q^2)}, U_2 = 1, U_3 \geq \frac{q^{\lambda_a} + 1}{q^{\lambda_a - 1} + 1}.$$

It follows that

$$\frac{\chi(1)}{\chi'(1)} \geq \frac{q^n + 1}{q^{n-1} + 1} \cdot \frac{(q^{\lambda_a} - q)(q^{\lambda_a} - q^2)}{(q^{\lambda_a - 1} - q)(q^{\lambda_a - 1} - q^2)} \cdot \frac{q^{\lambda_a} + 1}{q^{\lambda_a - 1} + 1} > q^4,$$

so $\chi(1) > q^{4n-10}$ as required.

5) Similarly, suppose that $(\lambda_1, \mu_1) = (0, 1)$ but $\mu \neq (1, 2, ..., b)$. Then there exists $j \geq 2$ such that $\mu_j \geq \mu_{j-1} + 2$. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n - 1$, with $\lambda' = \lambda$ and $\mu' = (\mu_1, ..., \mu_{j-1}, \mu_j - 1, \mu_{j+1}, ..., \mu_b)$. By the induction hypothesis, $\chi'(1) > q^{4n-14}$. Set

$$V_1 = \prod_{j' < j} \frac{q^{\mu_j} - q^{\mu_{j'}}}{q^{\mu_j - 1} - q^{\mu_{j'}}}, V_2 = \prod_{j' > j} \frac{q^{\mu_{j'}} - q^{\mu_j}}{q^{\mu_{j'}} - q^{\mu_j - 1}}, V_3 = \prod_i \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i} + q^{\mu_j - 1}}.$$

If $n - \mu_j \geq 3$ then

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n + 1)(q^{n-1} - 1)}{q^{2\mu_j} - 1} V_1 V_2 V_3 > \frac{(q^n + 1)(q^{n-1} - 1)}{q^{2\mu_j} - 1} \cdot \frac{q^{\mu_j} - q}{q^{\mu_j - 1} - q} \cdot \frac{1}{2} \cdot \frac{q^{\mu_j} + 1}{q^{\mu_j - 1} + 1} >$$

$$> \frac{(q^n + 1)(q^{n-1} - 1)q^2}{2(q^{2\mu_j} - 1)} > q^4,$$

so $\chi(1) > q^{4n-10}$ as desired. If $n - \mu_j = 2$ then $j = b$ and therefore $V_2 = 1$. We have $\chi(1)/\chi'(1) > (q^n + 1)(q^{n-1} - 1)q^2/(q^{2\mu_j} - 1) > q^4$ and we are done. Now suppose that $n - \mu_j \leq 1$. Then $n - \mu_j = 1$, $a - b = 2$, and

$$\binom{\lambda}{\mu} = \begin{pmatrix} 0 & 1 & ... & a-2 & a-1 \\ 1 & 2 & ... & b-1 & n-1 \end{pmatrix}.$$

As $b \geq 2$, we have $a \geq 4$. It follows that $V_1 > q$, $V_3 \geq (q^{n-1} + 1)/(q^{n-5} + 1) > q^3$. Hence $\chi(1)/\chi'(1) > q^4$ and we are done again.

6) Here we suppose that $\mu_1 = 0$ and $\lambda = (1, 2, ..., a)$. First we consider the case where $\mu \neq (0, 1, ..., b - 1)$. Then there exists an index $j \geq 2$ such that $\mu_j \geq \mu_{j-1} + 2$. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n - 1$, with $\lambda' = \lambda$ and $\mu' = (\mu_1, ..., \mu_{j-1}, \mu_j - 1, \mu_{j+1}, ..., \mu_b)$. Since $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$, by the induction hypothesis, we have $\chi'(1) > q^{4n-14}$. Observe that $n - \mu_j \geq (a - b + 2)^2/4 \geq 4$. So we have

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n + 1)(q^{n-1} - 1)}{q^{2\mu_j} - 1} \cdot \prod_{j' < j} \frac{q^{\mu_j} - q^{\mu_{j'}}}{q^{\mu_j - 1} - q^{\mu_{j'}}} \cdot \prod_{j' > j} \frac{q^{\mu_{j'}} - q^{\mu_j}}{q^{\mu_{j'}} - q^{\mu_j - 1}} \cdot \prod_i \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i} + q^{\mu_j - 1}} >$$

$$> \frac{(q^n+1)(q^{n-1}-1)}{2(q^{2\mu_j}-1)} > q^4,$$

and hence $\chi(1) > q^{4n-10}$.

Next we consider the case where $\mu = (0, 1, ..., b-1)$. Observe that $n = a+(a-b)^2/4$. If $b = 1$ then $n = (a+1)^2/4$ and $\chi(1) > (q^n+1)f(a+1)$ where f is the function defined in formula (3.4). Since $n \geq 6$, it follows that $a \geq 7$. We have already proved that $f(a+1) > q^{3n-10}$ for $a \geq 5$. Therefore $\chi(1) > q^{4n-10}$ as required. Now we can assume $b \geq 2$. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = (1, 2, ..., a-1)$ and $\mu' = (0, 1, ..., b-2)$. Since $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$, again by the induction hypothesis, $\chi'(1) > q^{4n-14}$. Furthermore,

$$\frac{\chi(1)}{\chi'(1)} = \frac{q^{a+b-2}(q^n+1)(q^{n-1}-1)}{(q^a-1)(q^{b-1}+1)} > q^{2n-3} > q^4.$$

Consequently, $\chi(1) > q^{4n-10}$.

7) Finally, we suppose that $\lambda_1 = 0$ and $\mu = (1, 2, ..., b)$. First we consider the case where $\lambda \neq (0, 1, ..., a-1)$. Then there exists $i \geq 2$ such that $\lambda_i \geq \lambda_{i-1}+2$. Choose i to be smallest possible. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = (\lambda_1, ..., \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, ..., \lambda_a)$ and $\mu' = \mu$. If $\binom{\lambda'}{\mu'} \in \mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu} = \binom{0\ 2\ n-1}{1}$ and by checking directly we have $\chi(1) > q^{4n-10}$. Therefore we can suppose that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. In other words, by the induction hypothesis, $\chi'(1) > q^{4n-14}$. Remark that $n - \lambda_i \geq 0$. Set

$$W_1 = \prod_{i'<i} \frac{q^{\lambda_i} - q^{\lambda_{i'}}}{q^{\lambda_i - 1} - q^{\lambda_{i'}}}, W_2 = \prod_{i'>i} \frac{q^{\lambda_{i'}} - q^{\lambda_i}}{q^{\lambda_{i'}} - q^{\lambda_i - 1}}, W_3 = \prod_{j} \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i - 1} + q^{\mu_j}}.$$

Note that $W_1 \geq (q^{\lambda_i}-1)/(q^{\lambda_i - 1}-1) > q$, $W_2 > 1/2$ and $W_3 > 1$. Therefore, if $n - \lambda_i \geq 3$,

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n+1)(q^{n-1}-1)}{q^{2\lambda_i}-1} W_1 W_2 W_3 > \frac{(q^n+1)(q^{n-1}-1)q}{2(q^{2\lambda_i}-1)} > q^4,$$

so $\chi(1) > q^{4n-10}$. If $1 \leq n - \lambda_i \leq 2$ then either $i = a$ or

$$\binom{\lambda}{\mu} = \begin{pmatrix} 0 & ... & n-4 & n-2 & n-1 \\ & 1 & ... & n-3 & \end{pmatrix}.$$

In the former case, $W_2 = 1$ and we have

$$\frac{\chi(1)}{\chi'(1)} \geq \frac{(q^n+1)(q^{n-1}-1)}{q^{2\lambda_i}-1} \cdot \frac{(q^{\lambda_i}-1)(q^{\lambda_i}-q)}{(q^{\lambda_i - 1}-1)(q^{\lambda_i - 1}-q)} \cdot \frac{q^{\lambda_i}+q}{q^{\lambda_i - 1}+q} >$$

$$> \frac{(q^n+1)(q^{n-1}-1)}{(q^{n-1}+1)(q^{n-2}-1)} \cdot \frac{q^{2\lambda_i}-q^2}{q^{2\lambda_i - 2}-q^2} > q^4,$$

and therefore we are done. In the latter case, since $n \geq 6$, $W_1 \geq q^4$ and we are done also. If $n = \lambda_i$, then $i = a$, $a - b = 2$ and $\lambda = (0, 1, ..., a-2, n)$. Since $\binom{\lambda}{\mu}$ is not in $\mathcal{L}_n$, $a \geq 4$. Then we have

$$\frac{\chi(1)}{\chi'(1)} = \prod_{i=1}^{a-2} \frac{q^n - q^i}{q^{n-1} - q^i} \cdot \prod_{j=1}^{a-2} \frac{q^n + q^j}{q^{n-1} + q^j} > q^{2(a-2)} \geq q^4,$$

so $\chi(1) > q^{4n-10}$.

The last configuration we have to handle is that $\lambda = (0, 1, ..., a-1)$ and $\mu = (1, 2, ..., b)$. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = (0, 1, ..., a-2)$ and $\mu' = (1, 2, ..., b-1)$ (if $b = 1$, μ' is just empty). By the induction hypothesis, $\chi'(1) > q^{4n-14}$. Furthermore,

$$\frac{\chi(1)}{\chi'(1)} = \frac{q^{a+b-2}(q^n+1)(q^{n-1}-1)}{(q^a-1)(q^{b-1}+1)} > q^{2n-3} > q^4.$$

Consequently, $\chi(1) > q^{4n-10}$. $\square$

3.3. **Unipotent Characters of $P(CO_{2n}^{+}(q)^0)$.** Proposition 7.2 in [TZ1] shows that the projective conformal orthogonal group of type +, $P(CO_{2n}^{+}(q)^0)$, has a unique unipotent character of minimal degree $(q^n-1)(q^{n-1}+q)/(q^2-1)$ and any other nontrivial unipotent character has degree greater than q^{2n-2}. We mimic its proof and get Proposition 3.4, which classifies unipotent characters of degrees up to q^{4n-10}.

From [C1, p. 471, 472], we know that the unipotent characters of G are labeled by symbols of the form

$$\binom{\lambda}{\mu} = \begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 & \ldots & \lambda_a \\ & \mu_1 & \mu_2 & \ldots & \mu_b \end{pmatrix},$$

where $0 \leq \lambda_1 < \lambda_2 < ... < \lambda_a$, $0 \leq \mu_1 < \mu_2 < ... < \mu_b$, $a - b \equiv 0 (\mathrm{mod}\ 4)$, $(\lambda_1, \mu_1) \neq (0, 0)$, and

$$\sum_i \lambda_i + \sum_j \mu_j - \left[\left(\frac{a+b-1}{2}\right)^2\right] = n.$$

The integer n is called the rank of the symbol $\binom{\lambda}{\mu}$. The degree of the unipotent character $\chi^{\lambda,\mu}$ corresponding to the symbol $\binom{\lambda}{\mu}$ is equal to

$$\frac{(q^n-1)\prod_{i=1}^{n-1}(q^{2i}-1)\prod_{i'<i}(q^{\lambda_i}-q^{\lambda_{i'}})\prod_{j'<j}(q^{\mu_j}-q^{\mu_{j'}})\prod_{i,j}(q^{\lambda_i}+q^{\mu_j})}{2^c q^{\binom{a+b-2}{2}+\binom{a+b-4}{2}+\cdots}\prod_i\prod_{k=1}^{\lambda_i}(q^{2k}-1)\prod_j\prod_{k=1}^{\mu_j}(q^{2k}-1)}.$$

Here $c = (a+b-2)/2$ if $\lambda \neq \mu$, and $c = a$ if $\lambda = \mu$. In the former case, $\binom{\lambda}{\mu}$ and $\binom{\mu}{\lambda}$ determine the same unipotent character. In the latter case, there are two unipotent characters with the same symbols.

Proposition 3.4. *Let* $G^* = (D_n)_{ad}(q) = P(CO^+_{2n}(q)^0)$*. Suppose that* $n \geq 5$*,* $(n,q) \neq (5,2)$*, and* $\chi \in \mathrm{Irr}(G^*)$ *is unipotent. Then either* χ *is one of characters labeled by* $\binom{n}{0}$*,* $\binom{n-1}{1}$*,* $\binom{0\;1}{1\;n}$ *with degrees* 1*,* $(q^n-1)(q^{n-1}+q)/(q^2-1)$*,* $(q^{2n}-q^2)/(q^2-1)$*, respectively, or* $\chi(1) > q^{4n-10}$*. Furthermore, when* $(n,q) = (5,2)$*,* G^* *has one more character of degree* 868 *(corresponding to the symbol* $\binom{0\;1\;2\;4}{-}$*), which is less than* q^{4n-10}*.*

Proof. When $n = 5$, Proposition can be verified directly by using [CH, Table 30]. Denote $\mathcal{L}_n = \{\{(n),(0)\},\{(n-1),(1)\},\{(0,1),(1,n)\}\}$. We will prove by induction on $n \geq 5$ that if $\chi = \chi^{\lambda,\mu}$ and $\{\lambda,\mu\} \notin \mathcal{L}_n$, then $\chi(1) > q^{4n-10}$, provided $(n,q) \neq (5,2)$. The rest of the proof establishes the induction step for $n \geq 6$.

1) At this point we suppose that $(\lambda_1,\mu_1) \neq (1,0)$ and $\lambda_1 \geq 1$ (eventually b may be zero). Consider the unipotent character χ' labeled by the symbol $\binom{\lambda'}{\mu'}$ of rank $n-1$, where $\lambda' = (\lambda_1-1,\lambda_2,...,\lambda_a)$ and $\mu' = \mu$. If $\{\lambda,\mu\} \in \mathcal{L}_{n-1}$ then $\binom{\lambda}{\mu} = \binom{2}{n-2}$ or $\binom{2\;n-1}{0\;1}$ and therefore $\chi(1) > q^{4n-10}$. So we can suppose that $\binom{\lambda'}{\mu'}$ is not in $\mathcal{L}_{n-1}$. By induction hypothesis, $\chi'(1) > q^{4(n-1)-10}$. The condition $n = \sum_i \lambda_i + \sum_j \mu_j - \frac{(a+b)^2-2(a+b)}{4} \geq a\lambda_1 + \frac{(a-b)^2}{4}$ implies that $n - \lambda_1 \geq 0$. If $\lambda_1 \leq n-4$ then

$$\frac{\chi(1)}{\chi'(1)} \geq \frac{(q^n-1)(q^{n-1}+1)}{2(q^{2\lambda_1}-1)} \cdot \prod_{i=2}^{a} \frac{q^{\lambda_i}-q^{\lambda_1}}{q^{\lambda_i}-q^{\lambda_1-1}} \cdot \prod_{j=1}^{b} \frac{q^{\lambda_1}+q^{\mu_j}}{q^{\lambda_1-1}+q^{\mu_j}} >$$

$$> \frac{(q^n-1)(q^{n-1}+1)}{4(q^{2\lambda_1}-1)} \geq \frac{(q^n-1)(q^{n-1}+1)}{4(q^{2n-8}-1)} > q^4,$$

so $\chi(1) > q^{4n-10}$. If $n - \lambda_1 \leq 3$ then $a = 1$ and $(a-b)^2/4 \leq 3$. Since $a - b \equiv 0 (\mathrm{mod}\ 4)$, $a = b = 1$. Note that $\{\lambda,\mu\}$ is not in $\mathcal{L}_n$. So $\binom{\lambda}{\mu} = \binom{n-2}{2}$ or $\binom{n-3}{3}$. Checking directly, we have $\chi(1) > q^{4n-10}$ as required.

The case $(\lambda_1,\mu_1) \neq (0,1)$ and $\mu_1 \geq 1$ can be reduced to 1) by interchanging λ and μ.

2) Now we consider the case where $b = 0$ and $\lambda_1 = 0$. Then $a \equiv 0 (\mathrm{mod}\ 4)$. First we suppose that $\lambda \neq (0,1,...,a-1)$. Then there exists $i \geq 2$ such that $\lambda_i - \lambda_{i-1} \geq 2$. We choose i to be smallest possible. Consider the character χ' corresponding to the symbol $\binom{\lambda'}{-}$, where $\lambda' = (...,\lambda_{i-1},\lambda_i-1,\lambda_{i+1},...)$. Note that $n - \lambda_i \geq (a-2)^2/4$. Therefore, $\chi(1)/\chi'(1) > (q^n+1)(q^{n-1}-1)/2(q^{2\lambda_i}-1) > q^4$ if $a \geq 8$. It is obvious that $\binom{\lambda'}{-} \notin \mathcal{L}_{n-1}$. If $(n,q) = (6,2)$ and $\lambda' = (0,1,2,5)$ or $(0,1,3,4)$, we can check directly that $\chi(1) > q^{4n-10}$. Otherwise, the induction hypothesis $\chi'(1) > q^{4n-14}$ implies that $\chi(1) > q^{4n-10}$. If $a = 4$ and $n - \lambda_i \geq 3$, then again $\chi(1)/\chi'(1) > q^4$, so $\chi(1) > q^{4n-10}$. If $a = 4$ and $n - \lambda_i \leq 2$, then $i = a$ and $\lambda = (0,1,2,n-1)$. Direct computation shows that $\chi(1) > q^{4n-10}$ and we are done. Next we suppose that $\lambda = (0,1,...,a-1)$. Then $\chi(1) = (q^n-1)f(a)$ and $n = a^2/4$, where $f(a)$ is the function defined in (3.4).

Note that $n \geq 6$, so $a \geq 8$. We already showed that $f(a) > q^{3n}$ for $a \geq 6$. Hence, $\chi(1) > q^{4n-10}$ as desired.

Again, the case $a = 0$ and $\mu_1 = 0$ can be reduced to 2) by interchanging λ and μ. Therefore, it remains to consider the case where $a, b \geq 1$ and $(\lambda_1, \mu_1) = (0, 1)$ or $(1, 0)$. By a similar reason as above, it is enough to suppose that $(\lambda_1, \mu_1) = (1, 0)$.

3) Here we suppose that $\lambda \neq (1, 2, ..., a)$. Then there exists $i \geq 2$ such that $\lambda_i \geq \lambda_{i-1} + 2$. Choose i to be smallest possible. Consider χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = (\lambda_1, ..., \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, ..., \lambda_a)$ and $\mu' = \mu$. By induction hypothesis, $\chi'(1) > q^{4n-14}$. Set

$$U_1 = \prod_{i'<i} \frac{q^{\lambda_i} - q^{\lambda_{i'}}}{q^{\lambda_i - 1} - q^{\lambda_{i'}}}, U_2 = \prod_{i'>i} \frac{q^{\lambda_{i'}} - q^{\lambda_i}}{q^{\lambda_{i'}} - q^{\lambda_i - 1}}, U_3 = \prod_j \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i - 1} + q^{\mu_j}}.$$

If $n - \lambda_i \geq 2$ then

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n-1)(q^{n-1}+1)}{q^{2\lambda_i}-1} U_1 U_2 U_3 > \frac{(q^n-1)(q^{n-1}+1)}{q^{2\lambda_i}-1} \cdot \frac{q^{\lambda_i} - q}{q^{\lambda_i - 1} - q} \cdot \frac{1}{2} \cdot \frac{q^{\lambda_i}+1}{q^{\lambda_i-1}+1} >$$
$$> \frac{(q^n-1)(q^{n-1}+1)q^2}{2(q^{2\lambda_i}-1)} \geq \frac{(q^n-1)(q^{n-1}+1)q^2}{2(q^{2n-4}-1)} > q^4,$$

so $\chi(1) > q^{4n-10}$. If $n - \lambda_i \leq 1$ then $i = a = b$. Note that $n - \lambda_i \geq 0$. If $n - \lambda_i = 1$ then

$$\binom{\lambda}{\mu} = \begin{pmatrix} 1 & 2 & ... & a-1 & n-1 \\ & 0 & 1 & ... & a-2 & a \end{pmatrix}.$$

The case $a = 2$ can be checked directly. So we can suppose $a \geq 3$. Then

$$U_1 \geq \frac{(q^{n-1}-q)(q^{n-1}-q^2)}{(q^{n-2}-q)(q^{n-2}-q^2)}, U_2 = 1, U_3 \geq \frac{q^{n-1}+1}{q^{n-2}+1}.$$

It follows that

$$\frac{\chi(1)}{\chi'(1)} \geq \frac{q^n-1}{q^{n-1}-1} \cdot \frac{(q^{n-1}-q)(q^{n-1}-q^2)}{(q^{n-2}-q)(q^{n-2}-q^2)} \cdot \frac{q^{n-1}+1}{q^{n-2}+1} > q^4,$$

so $\chi(1) > q^{4n-10}$ as required. If $n - \lambda_i = 0$ then

$$\binom{\lambda}{\mu} = \begin{pmatrix} & 1 & 2 & ... & a-1 & n \\ 0 & 1 & ... & a-2 & a-1 & \end{pmatrix}.$$

Since $\{\lambda, \mu\}$ is not in $\mathcal{L}_n$, $a \geq 3$. Then

$$\frac{\chi(1)}{\chi'(1)} \geq \frac{(q^n-1)(q^{n-1}+1)}{q^{2n}-1} \cdot \frac{(q^n-q)(q^n-q^2)}{(q^{n-1}-q)(q^{n-1}-q^2)} \cdot \frac{(q^n+1)(q^n+q)(q^n+q^2)}{(q^{n-1}+1)(q^{n-1}+q)(q^{n-1}+q^2)}$$
$$= \frac{(q^n-q)(q^n-q^2)}{(q^{n-1}-q)(q^{n-1}-q^2)} \cdot \frac{(q^n+q)(q^n+q^2)}{(q^{n-1}+q)(q^{n-1}+q^2)} > q^4,$$

and therefore $\chi(1) > q^{4n-10}$.

4) Finally, we suppose that $\mu_1 = 0$ and $\lambda = (1, 2, ..., a)$. First we consider the case where $\mu \neq (0, 1, ..., b-1)$. Then there exists an index $j \geq 2$ such that $\mu_j \geq \mu_{j-1} + 2$. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = \lambda$ and $\mu' = (\mu_1, ..., \mu_{j-1}, \mu_j - 1, \mu_{j+1}, ..., \mu_b)$. If $\{\lambda', \mu'\} \in \mathcal{L}_{n-1}$ then $\binom{\lambda'}{\mu'} = \binom{1\ n-1}{0\ 2}$ and hence $\chi(1) > q^{4n-10}$ by direct computation. Therefore we can assume $\{\lambda', \mu'\}$ is not in $\mathcal{L}_{n-1}$. By the induction hypothesis, $\chi'(1) > q^{4n-14}$. Set

$$V_1 = \prod_{j'<j} \frac{q^{\mu_j} - q^{\mu_{j'}}}{q^{\mu_j - 1} - q^{\lambda_{j'}}}, V_2 = \prod_{j'>j} \frac{q^{\mu_{j'}} - q^{\mu_j}}{q^{\mu_{j'}} - q^{\mu_j - 1}}, V_3 = \prod_i \frac{q^{\lambda_i} + q^{\mu_j}}{q^{\lambda_i} + q^{\mu_j - 1}}.$$

Remark that $V_1 > q, V_2 > 1/2, V_3 > 1$ and $n - \mu_j \geq 1$. If $n - \mu_j \geq 3$, we have

$$\frac{\chi(1)}{\chi'(1)} = \frac{(q^n-1)(q^{n-1}+1)}{q^{2\mu_j} - 1} V_1 V_2 V_3 > \frac{(q^n-1)(q^{n-1}+1)q}{2(q^{2\mu_j} - 1)} \geq \frac{(q^n-1)(q^{n-1}+1)q}{2(q^{2n-6} - 1)} > q^4,$$

so $\chi(1) > q^{4n-10}$. If $n - \mu_j = 2$ then $j = b$ and $V_2 = 1$. We still have $\chi(1)/\chi'(1) > (q^n-1)(q^{n-1}+1)q/(q^{2n-4}-1) > q^4$ and we are done. The last case we need to handle is $n - \mu_j = 1$. Note that $j = b$ and $V_2 = 1$ in this case. If $b \geq 3$ then

$$V_1 \geq \frac{(q^{n-1}-1)(q^{n-1}-q)}{(q^{n-2}-1)(q^{n-2}-q)} > \frac{q(q^{n-1}-q)}{(q^{n-2}-q)}, V_3 \geq \frac{q^{n-1}+q}{q^{n-2}+q}.$$

It follows that

$$\frac{\chi(1)}{\chi'(1)} > \frac{(q^n-1)(q^{n-1}+1)}{q^{2n}-1} \cdot \frac{q(q^{n-1}-q)}{(q^{n-2}-q)} \cdot \frac{q^{n-1}+q}{q^{n-2}+q} > q^4,$$

so $\chi(1) > q^{4n-10}$. If $b \leq 2$ then $a = b = 2$ and $\binom{\lambda}{\mu} = \binom{1\ 2}{0\ n-1}$, hence $\chi(1) > q^{4n-10}$.

Next we consider the case where $\mu = (0, 1, ..., b-1)$. Consider the unipotent character χ' labeled by $\binom{\lambda'}{\mu'}$ of rank $n-1$, with $\lambda' = (1, 2, ..., a-1)$ and $\mu' = (0, 1, ..., b-2)$ (if $a = 1$, λ' is just empty; if $b = 1$, μ' is just empty). It is obvious that $\{\lambda', \mu'\}$ is not in $\mathcal{L}_{n-1}$ and therefore $\chi'(1) > q^{4n-14}$ by the induction hypothesis. Furthermore,

$$\frac{\chi(1)}{\chi'(1)} = \frac{q^{a+b-2}(q^n-1)(q^{n-1}+1)}{(q^a-1)(q^{b-1}+1)} > q^{2n-3} > q^4.$$

Consequently, $\chi(1) > q^{4n-10}$. □

4. Odd characteristic symplectic groups

The low-dimensional unipotent characters of $G = Sp_{2n}(q)$, q odd (up to degree $(q^{2n-2}-1)(q^{2n}-1)(q^{n-2}-1)(q^{n-2}-q^3)/2\ (q^2-1)(q^4-1)(q^3+1))$ are already classified in Proposition 3.1. Therefore, Theorem 1.1 will be proved if we can classify non-unipotent characters of G of degrees up to $q^{4n-10}(q^n-1)/2$. We know that the dual group of G is $G^* = SO_{2n+1}(q)$ and every non-unipotent character $\chi \in \mathrm{Irr}(G)$ is parameterized by a pair $((s), \psi)$ where (s) is a nontrivial geometric conjugacy class of a semi-simple element $s \in G^*$ and ψ is an irreducible unipotent character of the centralizer $C := C_{G^*}(s)$. Moreover, $\chi(1) = (G^* : C)_{p'}\psi(1)$.

Proposition 4.1. *Let $G = Sp_{2n}(q)$ where $n \geq 6$ and q is an odd prime power. Suppose that $\alpha, \alpha_1, \alpha_2 = \pm$ and $\chi \in \mathrm{Irr}(G)$ is not unipotent of degree less than $q^{4n-10}(q^n-1)/2$. Then one of the following holds:*

1) *$\chi(1) = (q^{2n-2}-1)(q^{2n}-1)/|A|_{p'}$, where $A = GL_2^{\alpha}(q)$, $\mathbb{Z}_{q^2-\alpha}$, or $\mathbb{Z}_{q-\alpha_1} \times \mathbb{Z}_{q-\alpha_2}$, and χ is parameterized by $((s), 1_C)$, where $C \simeq SO_{2n-3}(q) \times A$ or $\chi(1) = q(q^{2n-2}-1)(q^{2n}-1)/(q-\alpha)(q^2-1)$, and χ is parameterized by $((s), 1_{SO_{2n-3}(q)} \otimes \lambda)$, where $C \simeq SO_{2n-3}(q) \times GL_2^{\alpha}(q)$ and λ is the unique nontrivial unipotent character of degree q of $GL_2^{\alpha}(q)$.*
2) *$\chi(1) = (q^{n-1}+\alpha_1)(q^{2n}-1)/2(q-\alpha_2)$, and χ is parameterized by $((s), \lambda \otimes 1_{\mathbb{Z}_{q-\alpha_2}})$, where $C \simeq O_{2n-2}^{\alpha_1}(q) \times \mathbb{Z}_{q-\alpha_2}$ and λ is one of two extensions of $1_{SO_{2n-2}^{\alpha_1}(q)}$ to $O_{2n-2}^{\alpha_1}(q)$.*
3) *$\chi(1) = (q^{2n}-1)\omega(1)/(q-\alpha)$, and χ is parameterized by $((s), \omega \otimes 1_{\mathbb{Z}_{q-\alpha}})$, where $C \simeq SO_{2n-1}(q) \times \mathbb{Z}_{q-\alpha}$ and ω is one of unipotent characters of $SO_{2n-1}(q)$ of degrees 1, $(q^{n-1}-1)(q^{n-1}-q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}+q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}-q)/2(q-1)$, or $(q^{n-1}-1)(q^{n-1}+q)/2(q-1)$.*
4) *$\chi(1) = (q^{2n-2}-1)(q^{2n}-1)/2(q-\alpha_1)(q-\alpha_2)$, and χ is parameterized by $((s), \lambda \otimes 1_{SO_{2n-3}(q) \times \mathbb{Z}_{q-\alpha_2}})$, where $C \simeq O_2^{\alpha_1}(q) \times SO_{2n-3}(q) \times \mathbb{Z}_{q-\alpha_2}$ and λ is one of two extensions of $1_{SO_2^{\alpha_1}(q)}$ to $O_2^{\alpha_1}(q)$.*
5) *$\chi(1) = (q^{2n}-1)\omega(1)/2(q-\alpha)$, and χ is parameterized by $((s), \omega \otimes \lambda)$, where $C \simeq SO_{2n-1}(q) \times O_2^{\alpha}(q)$, ω is one of unipotent characters of $SO_{2n-1}(q)$ of degrees 1, $(q^{n-1}-1)(q^{n-1}-q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}+q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}-q)/2(q-1)$, or $(q^{n-1}-1)(q^{n-1}+q)/2(q-1)$, and λ is one of two extensions of $1_{SO_2^{\alpha}(q)}$ to $O_2^{\alpha}(q)$.*
6) *$\chi(1) = (q^n+1)\lambda(1)/2$, and χ is parameterized by $((s), \lambda)$, where $C \simeq O_{2n}^{+}(q)$ and λ is an extension of one of three unipotent characters of $SO_{2n}^{+}(q)$ of degrees 1, $(q^n-1)(q^{n-1}+q)/(q^2-1)$ or $(q^{2n}-q^2)/(q^2-1)$ to $O_{2n}^{+}(q)$.*
7) *$\chi(1) = (q^n-1)\lambda(1)/2$, and χ is parameterized by $((s), \lambda)$, where $C \simeq O_{2n}^{-}(q)$ and λ is an extension of one of three unipotent characters of $SO_{2n}^{-}(q)$ of degrees 1, $(q^n+1)(q^{n-1}-q)/(q^2-1)$ or $(q^{2n}-q^2)/(q^2-1)$ to $O_{2n}^{-}(q)$.*
8) *$\chi(1) = (q^{n-1}+\alpha)(q^{2n}-1)\omega(1)/2(q^2-1)$, and χ is parameterized by $((s), \omega \otimes \lambda)$, where $C \simeq SO_3(q) \times O_{2n-2}^{\alpha}(q)$, $\omega = 1_{SO_3(q)}$ or the unique nontrivial unipotent character of degree q of $SO_3(q)$, and λ is one of two extensions of $1_{SO_{2n-2}^{\alpha}(q)}$ to $O_{2n-2}^{\alpha}(q)$.*
9) *$\chi(1) = (q^{2n-2}-1)(q^{2n}-1)\lambda(1)/2(q^2-1)(q^2-\alpha)$, and χ is parameterized by $((s), 1_{SO_{2n-3}(q)} \otimes \lambda)$, where $C \simeq SO_{2n-3}(q) \times O_4^{\alpha}(q)$ and λ is one of two extensions of $1_{SO_4^{\alpha}(q)}$ or the unique unipotent character of degree q^2 of $SO_4^{\alpha}(q)$ to $O_4^{\alpha}(q)$ or λ is the unique unipotent character of degree $2q$ of $O_4^{+}(q)$.*

Proof. For short notation, we set $D(n) := (q^n-1)q^{4n-10}/2$. By Lemma 2.2, $C \simeq A\times B$ where $A \simeq \prod_{i=1}^t GL_{a_i}^{\alpha_i}(q^{k_i})$, $a_i, k_i \in \mathbb{N}$, $\alpha_i = \pm 1$, $\sum_{i=1}^t k_i a_i = n-m$ and $B = SO_{2k+1}(q)\times O^{\pm}_{2m-2k}(q)$, $0 \le k \le m \le n$. It is easy to show that $|B|_{p'} \le \prod_{i=1}^m (q^{2i}-1)$ and $|A|_{p'} \le \prod_{i=1}^{n-m}(q^i-(-1)^i)$. Therefore,

$$\frac{|G|_{p'}}{|C_{G^*}(s)|_{p'}} \ge \frac{(q^{2(m+1)}-1)\cdots(q^{2n}-1)}{(q+1)(q^2-1)\cdots(q^{n-m}-(-1)^{n-m})} := f(m,n).$$

1) Case $m \le n-3$. Since the function $f(x,n)$ is first increasing and then decreasing in the interval $[0,n]$, we have $f(m,n) \ge \min\{f(0,n), f(n-3,n)\}$. It is easy to check that both $f(0,n)$ and $f(n-3,n)$ are greater than $D(n)$.

2) Case $m = n-2$. First we suppose that $k = n-2$. Then $\chi(1) = \frac{(q^{2n-2}-1)(q^{2n}-1)}{|A|_{p'}}\psi(1)$, where $A = GL_2^{\alpha}(q)$, $GL_1^{\alpha}(q^2)$ or $GL_1^{\alpha_1}(q)\times GL_1^{\alpha_2}(q)$, $\alpha, \alpha_1, \alpha_2 := \pm 1$ and ψ is a unipotent character of $C = SO_{2n-3}(q)\times A$. Suppose that $\psi = \omega\otimes\lambda$ where ω, λ are unipotent characters of $SO_{2n-3}(q), A$, respectively. If ω is nontrivial then $\omega(1) \ge (q^{n-2}-1)(q^{n-2}-q)/2(q+1)$ by [TZ1, Proposition 5.1]. Then

$$\chi(1) \ge \frac{(q^{2n-2}-1)(q^{2n}-1)(q^{n-2}-1)(q^{n-2}-q)}{2(q+1)^2(q^2-1)} > D(n).$$

If ω is trivial then 1) holds since $GL_1^{\alpha}(q^2)$ or $GL_1^{\alpha_1}(q)\times GL_1^{\alpha_2}(q)$ has only one unipotent character, which is trivial, while $A = GL_2^{\alpha}(q)$ has two unipotent characters, the trivial one and the one of degree q.

Next, we consider $k = 0$ then

$$\chi(1) \ge \frac{(q^{n-2}-1)(q^{2n-2}-1)(q^{2n}-1)}{2(q+1)(q^2-1)} > D(n).$$

Finally, if $1 \le k \le n-3$, we have

$$\begin{aligned}\chi(1) &\ge \frac{(q^{2n-2}-1)(q^{2n}-1)}{(q+1)(q^2-1)}\cdot\frac{\prod_{i=k+1}^{n-2}(q^{2i}-1)\cdot(q^{n-2-k}-1)}{2\prod_{i=1}^{n-2-k}(q^{2i}-1)}\\ &> \frac{(q^{2n-2}-1)(q^{2n}-1)}{(q+1)(q^2-1)}\cdot\frac{1}{2}q^{2k(n-2-k)}(q^{n-2-k}-1)\\ &> \frac{(q^{2n-2}-1)(q^{2n}-1)}{(q+1)(q^2-1)}\cdot\frac{1}{2}q^{2(n-3)} > D(n).\end{aligned}$$

3) Case $m = n-1$. Assume $k = 0$. Then $\chi(1) = \frac{(q^{n-1}+\alpha_1)(q^{2n}-1)}{2(q-\alpha_2)}\psi(1)$, where ψ is a unipotent character of $C \simeq O_{2n-2}^{\alpha_1}(q)\times GL_1^{\alpha_2}(q) = (SO_{2n-2}^{\alpha_1}(q)\cdot 2)\times GL_1^{\alpha_2}(q)$, $\alpha_1, \alpha_2 = \pm 1$. Note that $GL_1^{\alpha_2}(q) \simeq \mathbb{Z}_{q-\alpha_2}$ has only one unipotent character, which is the trivial one. Suppose that ψ is an irreducible constituent of $(\omega)^{SO_{2n-2}^{\alpha_1}(q)\cdot 2}\otimes 1_{GL_1^{\alpha_2}(q)}$ where ω is a unipotent character of $SO_{2n-2}^{\alpha_1}(q)$. By Proposition 2.1, the unipotent characters

of $SO^{\pm}_{2n-2}(q)$ are of the form $\theta \circ f$, where θ runs over the unipotent characters of $P(CO^{\pm}_{2n-2}(q)^0)$ and f is the canonical homomorphism $f : SO^{\pm}_{2n-2} \to P(CO^{\pm}_{2n-2}(q)^0)$. In particular, degrees of the unipotent characters of $SO^{\pm}_{2n-2}(q)$ are the same as those of $P(CO^{\pm}_{2n-2}(q)^0)$. Therefore, by Propositions 7.1, 7.2 of [TZ1], if ω is nontrivial then $\omega(1) \geq (q^{n-1}+1)(q^{n-2}-q)/(q^2-1)$. Then

$$\chi(1) \geq \frac{(q^{n-1}+\alpha_1)(q^{2n}-1)}{2(q-\alpha_2)} \cdot \frac{(q^{n-1}+1)(q^{n-2}-q)}{q^2-1} > D(n).$$

Suppose that ω is trivial. Since $(1_{SO^{\alpha_1}_{2n-2}(q)})^{O^{\alpha_1}_{2n-2}(q)}$ is the sum of two different irreducible characters of degree 1, $O^{\alpha_1}_{2n-2}(q)$ has two unipotent characters of degree 1. So, in this case, $\psi = \lambda \otimes 1_{GL^{\alpha_2}_1(q)}$ where λ is one of two extensions of $1_{SO^{\alpha_1}_{2n-2}(q)}$ to $O^{\alpha_1}_{2n-2}(q)$ and therefore 2) holds.

Assume $k = n-1$. Then $\chi(1) = \frac{q^{2n}-1}{q-\alpha}\psi(1)$, $\alpha = \pm 1$, where ψ is a unipotent character of $SO_{2n-1}(q) \times GL^{\alpha}_1(q)$. Again, we have $\psi = \omega \otimes 1_{GL^{\alpha}_1(q)}$ where ω is a unipotent character of $SO_{2n-1}(q)$. By Corollary 3.2, ω is either one of characters of degrees 1, $(q^{n-1}-1)(q^{n-1}-q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}+q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}-q)/2(q-1)$, $(q^{n-1}-1)(q^{n-1}+q)/2(q-1)$ or $\omega(1) > q^{4n-12}$. In the former case, 3) holds. In the latter case, $\chi(1) > \frac{q^{2n}-1}{q-\alpha} q^{4n-12} > D(n)$.

Assume $k = 1$. Then

$$\chi(1) \geq (G^* : C)_{p'} \geq \frac{(q^{n-2}-1)(q^{2n-2}-1)(q^{2n}-1)}{2(q+1)(q^2-1)} > D(n).$$

Assume $k = n-2$. Then

$$\chi(1) = \frac{(q^{2n-2}-1)(q^{2n}-1)}{2(q-\alpha_1)(q-\alpha_2)}\psi(1),$$

where ψ is an irreducible unipotent character of $C \simeq SO_{2n-3}(q) \times O^{\alpha_1}_2(q) \times GL^{\alpha_2}_1(q)$, $\alpha_1, \alpha_2 = \pm$. Note that both $SO^{\alpha_1}_2(q))$ and $GL^{\alpha_2}_1(q)$ have only one unipotent character, which is the trivial one. Therefore, $\psi = \omega \otimes \lambda \otimes 1_{GL^{\alpha_2}_1(q)}$, where ω is a unipotent character of $SO_{2n-3}(q)$ and λ is one of two extensions of $1_{SO^{\alpha_1}_2(q)}$ to $O^{\alpha_1}_2(q)$. If ω is trivial then 4) holds. If ω is nontrivial then $\omega(1) \geq (q^{n-2}-1)(q^{n-2}-q)/2(q+1)$ by Proposition 5.1 of [TZ1]. Then,

$$\chi(1) \geq \frac{(q^{2n}-1)(q^{2n-2}-1)}{2(q+1)^2} \cdot \frac{(q^{n-2}-1)(q^{n-2}-q)}{2(q+1)} > D(n).$$

Finally, assume $2 \leq k \leq n-3$. We have

$$\chi(1) \geq \frac{q^{2n}-1}{q+1} \cdot \frac{(q^{2(k+1)}-1)\cdots(q^{2(n-1)}-1)}{2(q^2-1)\cdots(q^{2(n-1-k)}-1)} \cdot (q^{n-1-k}-1)$$

$$\geq \frac{q^{2n}-1}{q+1} \cdot \frac{1}{2} q^{2k(n-1-k)} \cdot (q^{n-1-k}-1) \geq \frac{q^{2n}-1}{q+1} \cdot \frac{1}{2} q^{4(n-3)} \cdot (q^2-1) > D(n).$$

4) Case $m = n$. Since χ is not unipotent, $k < n$. We have

$$\chi(1) = \frac{\prod_{i=k+1}^{n}(q^{2i}-1)\cdot(q^{n-k}+\alpha)}{2\prod_{i=1}^{n-k}(q^{2i}-1)}\psi(1),$$

where ψ is a unipotent character of $C \simeq SO_{2k+1}(q) \times O^{\alpha}_{2(n-k)}(q), \alpha = \pm$. So, $\chi(1) > \frac{1}{2}q^{2k(n-k)}(q^{n-k}-1)$.

Assume $k = n-1$. Then $\chi(1) = \frac{q^{2n}-1}{2(q-\alpha)}\psi(1)$. We have $\psi = \omega \otimes \lambda$, where ω is a unipotent characters of $SO_{2n-1}(q)$ and λ is one of two extensions of $1_{SO_2^{\alpha_1}(q)}$ to $O_2^{\alpha_1}(q)$. By Corollary 3.2, ω is either one of characters of degrees 1, $(q^{n-1}-1)(q^{n-1}-q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}+q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}-q)/2(q-1)$, $(q^{n-1}-1)(q^{n-1}+q)/2(q-1)$ or $\omega(1) > q^{4n-12}$. In the former case, 5) holds. In the latter case, $\chi(1) > \frac{q^{2n}-1}{2(q-\alpha)}q^{4n-12} > D(n)$.

Assume $k = 0$. Then $\chi(1) = \frac{q^n+\alpha}{2}\psi(1)$, where ψ is a unipotent character of $C \simeq O^{\alpha}_{2n}(q) = SO^{\alpha}_{2n}(q)\cdot 2$. Suppose that ψ is an irreducible constituent of ω^C, where ψ is a unipotent character of $SO^{\alpha}_{2n}(q)$. We have $\omega = \theta \circ f$ where f is the canonical homomorphism $f : SO^{\pm}_{2n}(q) \to P(CO^{\pm}_{2n}(q)^0)$ and θ is a unipotent character of $P(CO^{\pm}_{2n}(q)^0)$. There are two cases:

Case $\alpha = +$: By Proposition 3.4, θ is either one of characters of degrees 1, $(q^n-1)(q^{n-1}+q)/(q^2-1)$, $(q^{2n}-q^2)/(q^2-1)$, or $\theta(1) > q^{4n-10}$. In the latter case,

$$\chi(1) = \frac{q^n+1}{2}\psi(1) \geq \frac{q^n+1}{2}\omega(1) > \frac{q^n+1}{2}q^{4n-10} > D(n).$$

In the former case, by [ST], the character afforded by the rank 3 permutation module of $O^{+}_{2n}(q)$ on the set of all singular 1-spaces of $\mathbb{F}_q^{2n}$ is the sum of three unipotent characters of degrees 1, $(q^n-1)(q^{n-1}+q)/(q^2-1)$, and $(q^{2n}-q^2)/(q^2-1)$. Hence, 6) holds in this case.

Case $\alpha = -$: By Proposition 3.3, θ is either one of characters of degrees 1, $(q^n+1)(q^{n-1}-q)/(q^2-1)$, $(q^{2n}-q^2)/(q^2-1)$, or $\theta(1) > q^{4n-10}$. In the latter case,

$$\chi(1) = \frac{q^n-1}{2}\psi(1) \geq \frac{q^n-1}{2}\omega(1) > \frac{q^n-1}{2}q^{4n-10} = D(n).$$

In the former case, again by [ST], the character afforded by the rank 3 permutation module of $O^{-}_{2n}(q)$ on the set of all singular 1-spaces of $\mathbb{F}_q^{2n}$ is the sum of three unipotent characters of degrees 1, $(q^n+1)(q^{n-1}-q)/(q^2-1)$, and $(q^{2n}-q^2)/(q^2-1)$. Hence, 7) holds in this case.

Assume $k = 1$. Then $\chi(1) = \frac{(q^{2n}-1)(q^{n-1}+\alpha)}{2(q^2-1)}\psi(1)$, where ψ is a unipotent character of $C \simeq SO_3(q) \times O^{\alpha}_{2n-2}(q)$. Suppose that ψ is an irreducible constituent of $\omega \otimes \varphi^C$ where ω, φ are unipotent characters of $SO_3(q), SO^{\alpha}_{2n-2}(q)$, respectively. Let us consider the first case where φ is nontrivial. Then by Propositions 7.1, 7.2 of [TZ1], we have

$\varphi(1) \geq (q^{n-1}+1)(q^{n-2}-q)/(q^2-1)$. Then

$$\chi(1) \geq \frac{(q^{2n}-1)(q^{n-1}-1)(q^{n-1}+1)(q^{n-2}-q)}{2(q^2-1)^2} > D(n).$$

Now suppose $\varphi = 1_{SO^{\alpha}_{2n-2}(q)}$. Since $SO_3(q)$ has two unipotent characters, the trivial one and the unique nontrivial unipotent character of degree q, 8) holds in this case.

Assume $k = 2$. Then

$$\chi(1) = \frac{(q^{2n-2}-1)(q^{2n}-1)(q^{n-2}\pm 1)}{2(q^2-1)(q^4-1)}\psi(1) > D(n).$$

Assume $k = n-2$. Then

$$\chi(1) = \frac{(q^{2n-2}-1)(q^{2n}-1)}{2(q^2-1)(q^2-\alpha)}\psi(1),$$

where ψ is a unipotent character of $C \simeq SO_{2n-3}(q) \times O^{\alpha}_4(q), \alpha = \pm$. Suppose that ψ is an irreducible constituent of $\omega \otimes \varphi^{O^{\alpha}_4(q)}$, where ω, φ are unipotent characters of $SO_{2n-3}(q), SO^{\alpha}_4(q)$, respectively. If ω is nontrivial, by Proposition 5.1 of [TZ1], $\omega(1) \geq (q^{n-2}-1)(q^{n-2}-q)/2(q+1)$ and therefore

$$\chi(1) \geq \frac{(q^{2n-2}-1)(q^{2n}-1)(q^{n-2}-1)(q^{n-2}-q)}{4(q+1)(q^4-1)} > D(n).$$

Now we assume $\omega = 1_{SO_{2n-3}(q)}$. There are two cases:

Case $\alpha = +$: Let $V := M_{2\times 2}(\mathbb{F}_q)$ be the space of 2×2 matrices over $\mathbb{F}_q$. Then V is a vector space over $\mathbb{F}_q$ of rank 4. The determinant function $Q(M) = \det(M)$, $M \in V$ is a quadratic form on V with Witt index 2. So the group of linear transformations $V \to V$ preserving Q will be $O^+_4(q)$. Consider an action of group $GL_2(q) \times GL_2(q)$ on V by $\tau(A,B)(M) = A^{-1}MB$. We see that $\tau(A,B)$ preserves Q if and only if $\det(A) = \det(B)$, and $\tau(A,B)$ is the identity if and only if $A = B$ is a scalar matrix. Moreover, $\tau(A,B)$ has determinant $\det(A)^{-2}\det(B)^2$, which is 1 when $\det(A) = \det(B)$. Therefore, we have a homomorphism

$$\tau : (SL_2(q) \times SL_2(q)) \cdot \mathbb{Z}_2 \to SO^+_4(q),$$

where $\mathbb{Z}_2$ is generated by

$$U = \left(\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \right).$$

We have $\mathrm{Ker}(\rho) = \mathbb{Z}_2$, which is generated by $(-I,-I)$. Therefore, $((SL_2(q)\times SL_2(q))\cdot \mathbb{Z}_2)/\mathbb{Z}_2 \simeq 2\cdot(PSL_2(q)\times PSL_2(q))\cdot 2$ is a subgroup of $SO^+_4(q)$. Since $|2\cdot(PSL_2(q)\times PSL_2(q))\cdot 2| = |SO^+_4(q)| = q^2(q^2-1)^2$, $2\cdot(PSL_2(q)\times PSL_2(q))\cdot 2 \simeq SO^+_4(q)$. We denote by L_1, L_2 the first and second terms respectively in $PSL_2(q)\times PSL_2(q)$. Note that $P(CO^+_4(q)^0)$ as well as $SO^+_4(q)$ have four unipotent characters of degrees $1, q, q, q^2$. Since $PSL_2(q)$ has only one character of degree q which is denoted by μ_1 for L_1 and μ_2 for L_2 and U fixes L_1 and L_2, the unipotent characters of $SO^+_4(q)$ must be extensions of characters $1_{L_1\times L_2}, \mu_1\otimes 1_{L_2}, 1_{L_1}\otimes\mu_2$ and $\mu_1\otimes\mu_2$ which are considered as characters of $2\cdot(L_1\times L_2)$.

Consider the transformation $T: V\to V$ defined by $T(M)=M^T$, the transpose of M. It is clear that T preserves Q and therefore $T\in O_4^+(q)$. Moreover $\det(T)=-1$. So $O_4^+(q)=SO_4^+(q)\rtimes <T>$. We have $T^{-1}\tau(A,B)T(M)=B^TM(A^T)^{-1}$ for every $A,B\in GL_2(q)$. So T fixes $\mu_1\otimes\mu_2$ and maps one of $\{\mu_1\otimes 1_{L_2}, 1_{L_1}\otimes\mu_2\}$ to the other. In other words, the unipotent character of degree q^2 of $SO_4^+(q)$ has two extensions to $O_4^+(q)$ and the inductions of two unipotent characters of degree q of $SO_4^+(q)$ to $O_4^+(q)$ are equal and irreducible. So 9) holds in this case.

Case $\alpha=-$: Note that $SO_4^-(q)$ as well as $P(CO_4^-(q)^0)$ have two unipotent characters of degrees $1, q^2$. It is well known that $SO_4^-(q)\simeq PSL_2(q^2)\times 2_1$ and $(O_4^-(q)=(PSL_2(q^2)\times 2_1)\cdot 2_2\simeq 2_1\times(PSL_2(q^2)\cdot 2_2)$, where 2_2 acts trivially on 2_1. Since $PSL_2(q^2)$ has only one irreducible character of degree q^2 which is unipotent, the unipotent character of degree q^2 of $SO_4^-(q)$ is invariant in $O_4^-(q)$. Therefore, $O_4^-(q)$ has two unipotent characters of degree q^2. So 9) also holds in this case.

Lastly, we assume $3\le k\le n-3$. Then

$$\chi(1)\ge\frac{\prod_{i=k+1}^{n}(q^{2i}-1)\cdot(q^{n-k}\pm 1)}{2\prod_{i=1}^{n-k}(q^{2i}-1)}>$$

$$>\frac{1}{2}q^{2k(n-k)}(q^{n-k}\pm 1)\ge\frac{1}{2}q^{6(n-3)}(q^3-1)>D(n).$$

□

Counting semi-simple conjugacy classes in $\mathbf{G^*=SO_{2n+1}(q)}$: In Proposition 4.1 and its proof, we have not shown how to count the number of semi-simple conjugacy classes (s) in $SO_{2n+1}(q)$ for a certain $C:=C_{G^*}(s)$, which will imply the number of irreducible characters of $Sp_{2n}(q)$ at each degree. Actually, the way to count them is pretty similar in all the cases from 1) to 9) in Proposition 4.1. First, from the structure of $C_{G^*}(s)$, we know the form of the characteristic polynomial as well as the eigenvalues of s. In general, we have

$$\mathrm{Spec}(s)=\{\underbrace{1,...,1}_{2k+1},\underbrace{-1,...,-1}_{2(m-k)},\underbrace{...}_{2(n-m)}\}.$$

By [W1, (2.6)], if $-1\notin\mathrm{Spec}(s)$, there is exactly one $O_{2n+1}(q)$-conjugacy class of semi-simple elements (s) for a given $\mathrm{Spec}(s)$. This class is also an $SO_{2n+1}(q)$-conjugacy class since $(O_{2n+1}(q):C_{O_{2n+1}(q)}(s))=(SO_{2n+1}(q):C_{SO_{2n+1}(q)}(s))$ (see Lemma 2.2). The situation is a little bit different when $-1\in\mathrm{Spec}(s)$. In that case, again by [W1, (2.6)], there are exactly two $O_{2n+1}(q)$-conjugacy class of semi-simple elements (s) for a given $\mathrm{Spec}(s)$, in which $C_{O_{2n+1}(q)}(s)\simeq O_{2k+1}(q)\times O^{\pm}_{2(m-k)}(q)\times\prod_{i=1}^{t}GL_{a_i}^{\alpha_i}(q^{k_i})$ (+ for one class and − for the other). These classes are also $SO_{2n+1}(q)$-conjugacy class by the same reason as before. So, in order to count the number of semi-simple conjugacy classes (s) with a given centralizer C, we need to count the number of choices of $\mathrm{Spec}(s)$. This is demonstrated in Tables 2, 3.

We finish this section by two following Corollaries of Theorem 1.1.

Corollary 4.2. *Let χ be an irreducible complex character of $G = Sp_{2n}(q)$, where $n \geq 6$ and q is an odd prime power. Then $\chi(1) = 1$ (1 character), $(q^n \pm 1)/2$ (4 characters), $(q^n + \alpha_1)(q^n + \alpha_2 q)/2(q + \alpha_1\alpha_2)$ (4 characters) $(\alpha_{1,2} = \pm 1)$, $(q^{2n} - 1)/2(q \pm 1)$ (4 characters), $(q^{2n} - 1)/(q \pm 1)$ ($q - 2$ characters), or $\chi(1) \geq (q^{2n} - 1)(q^{n-1} - q)/2(q^2 - 1)$.*

Corollary 4.3. *Let χ be an irreducible complex character of $G = Sp_{2n}(q)$, where $n \geq 6$ and q is an odd prime power. Then $\chi(1) = 1$ (1 character), $(q^n \pm 1)/2$ (4 characters), $(q^n + \alpha_1)(q^n + \alpha_2 q)/2(q + \alpha_1\alpha_2)$ (4 characters) $(\alpha_{1,2} = \pm 1)$, $(q^{2n} - 1)/2(q \pm 1)$ (4 characters), $(q^{2n} - 1)/(q \pm 1)$ ($q - 2$ characters), $(q^{2n} - 1)(q^{n-1} \pm q)/2(q^2 - 1)$ (4 characters), $(q^{2n} - 1)(q^{n-1} \pm 1)/2(q^2 - 1)$ (4 characters), $(q^{2n} - 1)(q^{n-1} \pm 1)/2(q \pm 1)$ ($4q - 8$ characters), $(q^{2n} - q^2)(q^n \pm 1)/2(q^2 - 1)$ (4 characters), $q(q^{2n} - 1)(q^{n-1} \pm 1)/2(q^2 - 1)$ (4 characters), or $\chi(1) \geq (q^{2n} - 1)(q^{n-1} - 1)(q^{n-1} - q^2)/2(q^4 - 1)$.*

TABLE 1. Low-dimensional unipotent characters of $Sp_{2n}(q)$, $n \geq 6$, q odd. See §3.1 for the explanation of notation.

Characters	Symbols	Degrees
1_G	$\begin{pmatrix} n \\ - \end{pmatrix}$	1
χ_2	$\begin{pmatrix} 0\ 1\ n \\ - \end{pmatrix}$	$\dfrac{(q^n-1)(q^n-q)}{2(q+1)}$
χ_3	$\begin{pmatrix} 0\ 1 \\ n \end{pmatrix}$	$\dfrac{(q^n+1)(q^n+q)}{2(q+1)}$
χ_4	$\begin{pmatrix} 1\ n \\ 0 \end{pmatrix}$	$\dfrac{(q^n+1)(q^n-q)}{2(q-1)}$
χ_5	$\begin{pmatrix} 0\ n \\ 1 \end{pmatrix}$	$\dfrac{(q^n-1)(q^n+q)}{2(q-1)}$
χ_6	$\begin{pmatrix} 0\ 2\ n-1 \\ - \end{pmatrix}$	$\dfrac{(q^{2n}-1)(q^{n-1}-1)(q^{n-1}-q^2)}{2(q^4-1)}$
χ_7	$\begin{pmatrix} 0\ 2 \\ n-1 \end{pmatrix}$	$\dfrac{(q^{2n}-1)(q^{n-1}+1)(q^{n-1}+q^2)}{2(q^4-1)}$
χ_8	$\begin{pmatrix} 2\ n-1 \\ 0 \end{pmatrix}$	$\dfrac{(q^{2n}-1)(q^{n-1}+1)(q^{n-1}-q^2)}{2(q^2-1)^2}$
χ_9	$\begin{pmatrix} 0\ n-1 \\ 2 \end{pmatrix}$	$\dfrac{(q^{2n}-1)(q^{n-1}-1)(q^{n-1}+q^2)}{2(q^2-1)^2}$
χ_{10}	$\begin{pmatrix} 1\ n-1 \\ 1 \end{pmatrix}$	$\dfrac{q(q^{2n}-1)(q^{2n-2}-q^2)}{(q^2-1)^2}$
χ_{11}	$\begin{pmatrix} 0\ 1\ 2\ n \\ 1 \end{pmatrix}$	$\dfrac{(q^{2n}-q^2)(q^n-1)(q^n-q^2)}{2(q^4-1)}$
χ_{12}	$\begin{pmatrix} 0\ 1\ 2 \\ 1\ n \end{pmatrix}$	$\dfrac{(q^{2n}-q^2)(q^n+1)(q^n+q^2)}{2(q^4-1)}$
χ_{13}	$\begin{pmatrix} 1\ 2\ n \\ 0\ 1 \end{pmatrix}$	$\dfrac{(q^{2n}-q^2)(q^n+1)(q^n-q^2)}{2(q^2-1)^2}$
χ_{14}	$\begin{pmatrix} 0\ 1\ n \\ 1\ 2 \end{pmatrix}$	$\dfrac{(q^{2n}-q^2)(q^n-1)(q^n+q^2)}{2(q^2-1)^2}$

TABLE 2. Low-dimensional irreducible characters of $Sp_{2n}(q)$, $n \geq 6$, q odd. See §2.1 and Proposition 4.1 for the explanation of notation.

Spec(s)	$C_{G^\star}(s)$	$\psi(1)$	$\chi(1)$	♯ of characters
$\{1,...,1,\gamma^k,\gamma^{-k},\gamma^k,\gamma^{-k}\}$ $k \in T_1$	$SO_{2n-3}(q) \times GL_2(q)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{(q-1)(q^2-1)}$	$(q-3)/2$
		q	$\dfrac{q(q^{2n-2}-1)(q^{2n}-1)}{(q-1)(q^2-1)}$	$(q-3)/2$
$\{1,...,1,\eta^k,\eta^{-k},\eta^k,\eta^{-k}\}$ $k \in T_2$	$SO_{2n-3}(q) \times GU_2(q)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{(q+1)(q^2-1)}$	$(q-1)/2$
		q	$\dfrac{q(q^{2n-2}-1)(q^{2n}-1)}{(q+1)(q^2-1)}$	$(q-1)/2$
$\{1,...,1,\theta^j,\theta^{qj},\theta^{-j},\theta^{-qj}\}$ $j \in R_2$	$SO_{2n-3}(q) \times GL_1(q^2)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{q^2-1}$	$(q-1)^2/4$
$\{1,...,1,\zeta^j,\zeta^{qj},\zeta^{-j},\zeta^{-qj}\}$ $j \in R_1$	$SO_{2n-3}(q) \times GU_1(q^2)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{q^2+1}$	$(q^2-1)/4$
$\{1,...,1,\gamma^k,\gamma^{-k},\gamma^l,\gamma^{-l}\}$ $k,l \in T_1, k \neq l$	$SO_{2n-3}(q) \times GL_1(q) \times GL_1(q)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{(q-1)^2}$	$(q-3)(q-5)/8$
$\{1,...,1,\eta^k,\eta^{-k},\eta^l,\eta^{-l}\}$ $k,l \in T_2, k \neq l$	$SO_{2n-3}(q) \times GU_1(q) \times GU_1(q)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{(q+1)^2}$	$(q-1)(q-3)/8$
$\{1,...,1,\gamma^k,\gamma^{-k},\eta^l,\eta^{-l}\}$ $k \in T_1, l \in T_2$	$SO_{2n-3}(q) \times GL_1(q) \times GU_1(q)$	1	$\dfrac{(q^{2n-2}-1)(q^{2n}-1)}{q^2-1}$	$(q-1)(q-3)/4$
$\{1,-1,...,-1,\gamma^k,\gamma^{-k}\}$, $k \in T_1$	$O^\alpha_{2n-2}(q) \times GL_1(q)$	1	$\dfrac{(q^{2n}-1)(q^{n-1}+\alpha)}{2(q-1)}$	$2(q-3)$
$\{1,-1,...,-1,\eta^k,\eta^{-k}\}$, $k \in T_2$	$O^\alpha_{2n-2}(q) \times GU_1(q)$	1	$\dfrac{(q^{2n}-1)(q^{n-1}+\alpha)}{2(q+1)}$	$2(q-1)$
$\{1,...,1,\gamma^k,\gamma^{-k}\}$, $k \in T_1$	$SO_{2n-1}(q) \times GL_1(q)$	$(\star)$	$\dfrac{q^{2n}-1}{q-1}\psi(1)$	$(q-3)/2$
$\{1,...,1,\eta^k,\eta^{-k}\}$, $k \in T_2$	$SO_{2n-1}(q) \times GU_1(q)$	$(\star)$	$\dfrac{q^{2n}-1}{q+1}\psi(1)$	$(q-1)/2$

Table 3. Low-dimensional irreducible characters of $Sp_{2n}(q)$, $n \geq 6$, q odd. See §2.1 and Proposition 4.1 for the explanation of notation.

Spec(s)	$C_{G^\star}(s)$	$\psi(1)$	$\chi(1)$	♯ of chars
$\{1, ..., 1, -1, -1, \gamma^k, \gamma^{-k}\}$ $k \in T_1$	$SO_{2n-3}(q) \times O_2^\alpha(q)$ $\times GL_1(q)$	1	$\frac{(q^{2n-2}-1)(q^{2n}-1)}{2(q-\alpha)(q-1)}$	$2(q-3)$
$\{1, ..., 1, -1, -1, \eta^k, \eta^{-k}\}$ $k \in T_2$	$SO_{2n-3}(q) \times O_2^\alpha(q)$ $\times GU_1(q)$	1	$\frac{(q^{2n-2}-1)(q^{2n}-1)}{2(q-\alpha)(q+1)}$	$2(q-1)$
$\{1, ..., 1, -1, -1\}$	$SO_{2n-1}(q) \times O_2^\alpha(q)$	$(\star)$	$\frac{q^{2n}-1}{2(q-\alpha)}\psi(1)$	4
$\{1, -1, ..., -1\}$	$O_{2n}^\alpha(q)$	1	$(q^n+\alpha)/2$	4
		$\frac{(q^n-\alpha)(q^{n-1}+\alpha q)}{q^2-1}$	$\frac{(q^{2n}-1)(q^{n-1}+\alpha q)}{2(q^2-1)}$	4
		$\frac{q^{2n}-q^2}{q^2-1}$	$\frac{(q^{2n}-q^2)(q^n+\alpha)}{2(q^2-1)}$	4
$\{1, 1, 1, -1, ..., -1\}$	$SO_3(q) \times O_{2n-2}^\alpha(q)$	1	$\frac{(q^{2n}-1)(q^{n-1}+\alpha)}{2(q^2-1)}$	4
		q	$\frac{q(q^{2n}-1)(q^{n-1}+\alpha)}{2(q^2-1)}$	4
$\{1, ..., 1, -1, -1, -1, -1\}$	$SO_{2n-3}(q) \times O_4^\alpha(q)$	1	$\frac{(q^{2n-2}-1)(q^{2n}-1)}{2(q^2-1)(q^2-\alpha)}$	4
		q^2	$\frac{q^2(q^{2n-2}-1)(q^{2n}-1)}{2(q^2-1)(q^2-\alpha)}$	4
$\{1, ..., 1, -1, -1, -1, -1\}$	$SO_{2n-3}(q) \times O_4^+(q)$	$2q$	$\frac{q(q^{2n-2}-1)(q^{2n}-1)}{(q^2-1)^2}$	1

Where $(\star)$ is 1, $(q^{n-1}-1)(q^{n-1}-q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}+q)/2(q+1)$, $(q^{n-1}+1)(q^{n-1}-q)/2(q-1)$, or $(q^{n-1}-1)(q^{n-1}+q)/2(q-1)$.

5. Odd characteristic orthogonal groups in odd dimension

The aim of this section is to classify low-dimensional complex representations of groups $G = Spin_{2n+1}(q)$ where q is a power of an odd prime p. The dual group G^* is the projective conformal symplectic group $PCSp_{2n}(q)$, which is the quotient of $\widetilde{G} = CSp_{2n}(q)$ by its center $Z(\widetilde{G}) \simeq \mathbb{Z}_{q-1}$.

Proof of Theorem 1.2. If χ is unipotent, Theorem is done by Corollary 3.2. So we can assume χ is not unipotent such that $\chi(1) \leq q^{4n-8}$. Suppose that χ is parameterized by $((s^*), \psi)$, where s^* is a non-trivial semi-simple element in G^*. Consider an inverse image s of s^* in $\widetilde{G}$. Then (s) is a non-trivial conjugacy class of semi-simple elements in $\widetilde{G}$. Set $C^* = C_{G^*}(s^*)$ and $C = C_{\widetilde{G}}(s)$. Let g be any element in $\widetilde{G}$ such that the image of g in G^* belongs to C^*. Then $gsg^{-1} = \mu(g)s$ for some $\mu(g) \in \mathbb{Z}_{q-1}$. Therefore, $\tau(s) = \tau(gsg^{-1}) = \tau(\mu(g)s) = \mu(g)^2\tau(s)$. It follows that $\mu(g)$ is 1 or -1. Hence

$$(G^* : C^*)_{p'} \geq (\widetilde{G} : C)_{p'}/2.$$

Case 1: If $\tau(s)$ is not a square in $\mathbb{F}_q$, by Lemma 2.4, $C \simeq (Sp_m(q^2) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i})) \cdot \mathbb{Z}_{q-1}$, where $\alpha_i = \pm$, $\Sigma_{i=1}^{t} k_i a_i = n - m$. Therefore,

$$(\widetilde{G} : C)_{p'} \geq \frac{(q^2-1)\cdots(q^{2n}-1)}{(q^4-1)\cdots(q^{2m}-1)(q+1)(q^2-1)\cdots(q^{n-m}-(-1)^{n-m})}$$

$$\geq \begin{cases} \min\{\dfrac{(q^2-1)\cdots(q^{2n}-1)}{(q^4-1)\cdots(q^{2n}-1)}, \dfrac{(q^2-1)\cdots(q^{2n}-1)}{(q+1)(q^2-1)\cdots(q^n-(-1)^n)}\} & \text{if } 2 \mid n \\ \min\{\dfrac{(q^2-1)\cdots(q^{2n}-1)}{(q^4-1)\cdots(q^{2n-2}-1)(q+1)}, \dfrac{(q^2-1)\cdots(q^{2n}-1)}{(q+1)(q^2-1)\cdots(q^n-(-1)^n)}\} & \text{if } 2 \nmid n \end{cases}$$

$$= \begin{cases} \min\{(q^2-1)(q^6-1)\cdots(q^{2n-2}-1), (q-1)\cdots(q^n+(-1)^n)\} & \text{if } 2 \mid n \\ \min\{(q^2-1)(q^6-1)\cdots(q^{2n}-1)/(q+1), (q-1)\cdots(q^n+(-1)^n)\} & \text{if } 2 \nmid n, \end{cases}$$

which is greater than $2q^{4n-8}$. Therefore,

$$\chi(1) \geq (G^* : C^*)_{p'} \geq (\widetilde{G} : C)_{p'}/2 > q^{4n-8}.$$

Case 2: If $\tau(s)$ is a square in $\mathbb{F}_q$, multiplying s with a suitable scalar, we can assume that $\tau(s) = 1$ or in other words $s \in Sp_{2n}(q)$. By Lemma 2.4, $C \simeq (Sp_{2k}(q) \times Sp_{2(m-k)}(q) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i})) \cdot \mathbb{Z}_{q-1}$, where $\alpha_i = \pm$, $\Sigma_{i=1}^{t} k_i a_i = n - m$. It is easy to see that $|Sp_{2k}(q) \times Sp_{2(m-k)}(q)|_{p'} \leq \prod_{i=1}^{m}(q^{2i}-1)$ and $|\prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i})|_{p'} \leq \prod_{i=1}^{n-m}(q^i - (-1)^i)$. Therefore,

$$(\widetilde{G} : C)_{p'} \geq \frac{(q^{2(m+1)}-1)\cdots(q^{2n}-1)}{(q+1)(q^2-1)\cdots(q^{n-m}-(-1)^{n-m})} =: f(m,n).$$

1) Case $m \le n-2$. We have $(\widetilde{G}:C)_{p'} \geq \min\{f(0,n), f(n-2,n)\}$. Since both $f(0,n) = \prod_{i=1}^{n}(q^i + (-1)^i)$ and $f(n-2,n) = (q^{2n-2}-1)(q^{2n}-1)/(q+1)(q^2-1)$ are greater than $2q^{4n-8}$, $\chi(1) \geq (\widetilde{G}:C)_{p'}/2 > q^{4n-8}$.

2) Case $m = n-1$. If $1 \le k \le n-2$ then $(\widetilde{G}:C)_{p'} \geq (q^{2n-2}-1)(q^{2n}-1)/(q+1)(q^2-1)$. Again, $\chi(1) \geq (\widetilde{G}:C)_{p'}/2 > q^{4n-8}$. So $k=0$ or $n-1$. Modulo $Z(\widetilde{G})$, we can assume that $k = n-1$. There are two cases:

- $C = (Sp_{2n-2}(q) \times GL_1(q)) \cdot \mathbb{Z}_{q-1}$, which is happened when $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$, where $\pm 1 \neq \lambda \in \mathbb{F}_q^\times$. Note that there are $(q-3)/2$ choices for λ, namely, $\lambda = \gamma^k, k \in T_1$. For each such λ, there is exactly one semi-simple conjugacy class of such elements s in $Sp_{2n}(q)$ by [W1, (2.6)]. This conjugacy class is also the conjugacy class of s in $CSp_{2n}(q)$. Therefore, there is exactly one conjugacy class of semi-simple elements (s^*) in G^* such that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ for each $\lambda = \gamma^k, k \in T_1$.
- $C = (Sp_{2n-2}(q) \times GU_1(q)) \cdot \mathbb{Z}_{q-1}$, which is happened when $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$, where $\pm 1 \neq \lambda \in \mathbb{F}_{q^2}^\times$ and $\lambda^{-1} = \lambda^q$. Note that there are $(q-1)/2$ choices for λ, namely, $\lambda = \eta^k, k \in T_2$. Similarly as above, there is exactly one conjugacy class of semi-simple elements (s^*) in G^* such that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ for each $\lambda = \eta^k, k \in T_2$.

In two above situations, if $\{1, ..., 1, \lambda, \lambda^{-1}\} = \{\mu, ..., \mu, \mu\lambda, \mu\lambda^{-1}\}$ for some $\mu \in \mathbb{F}_q^\times$, then $\mu = 1$ since $n \geq 5$. Therefore, C is the complete inverse image of C^* in $\widetilde{G}$. In other words, $C^* = C/Z(\widetilde{G})$ and hence $(G^*:C^*)_{p'} = (\widetilde{G}:C)_{p'}$. Consider the canonical homomorphism $f: Sp_{2n-2}(q) \times GL_1^\alpha(q) \hookrightarrow C \to C^*$, $\alpha = \pm$, whose kernel is contained in the center of $Sp_{2n-2}(q) \times GL_1^\alpha(q)$ and image contains the commutator group of C^* since $Sp_{2n-2}(q) \times GL_1^\alpha(q)$ contains the commutator group of $C \simeq (Sp_{2n-2}(q) \times GL_1^\alpha(q)) \cdot \mathbb{Z}_{q-1}$. By Proposition 2.1, the unipotent characters of $Sp_{2n-2}(q) \times GL_1^\alpha(q)$ are of the form $\psi \circ f$, where ψ runs over the unipotent characters of C^*. In particular, ψ is trivial or $\psi(1) \geq (q^{n-1}-1)(q^{n-1}-q)/2(q+1)$ by [TZ1, Proposition 5.1]. In the latter case,

$$\chi(1) \geq \frac{q^{2n}-1}{q+1} \cdot \frac{(q^{n-1}-1)(q^{n-1}-q)}{2(q+1)} > q^{4n-8}.$$

Therefore, in this case, χ is one of $(q-3)/2$ characters of degree $(q^{2n}-1)/(q-1)$ or $(q-1)/2$ characters of degree $(q^{2n}-1)/(q+1)$.

3) Case $m = n$. Since (s^*) is non-trivial, we assume $1 \le k \le n-1$.

First, if $k = 1$ or $n-1$, modulo $Z(\widetilde{G})$, we may assume $\mathrm{Spec}(s) = \{-1, -1, 1, ..., 1\}$. Again, $C^* = C/Z(\widetilde{G})$ and $(G^*:C^*)_{p'} = (\widetilde{G}:C)_{p'}$. There is a unique possibility for (s^*) in this case. We have $\chi(1) = \frac{q^{2n}-1}{q^2-1}\psi(1)$ where ψ is a unipotent character of C^*. Applying Proposition 2.1 again for the canonical homomorphism $f: Sp_2(q) \times Sp_{2n-2}(q) \hookrightarrow C \to C^*$, we see that either ψ is trivial, or the unipotent character of degree q, or $\psi(1) \geq (q^{n-1}-1)(q^{n-1}-q)/2(q+1)$. In the last case,

$$\chi(1) \geq \frac{q^{2n}-1}{q^2-1} \cdot \frac{(q^{n-1}-1)(q^{n-1}-q)}{2(q+1)} > q^{4n-8}.$$

The first two cases explain why G has two characters of degrees $(q^{2n}-1)/(q^2-1)$ and $q(q^{2n}-1)/(q^2-1)$.

Next, if $k = 2$ or $n-2$, since $n \geq 5$, again one can show that $C^* = C/Z(\widetilde{G})$ and hence

$$\chi(1) \geq (\widetilde{G} : C)_{p'} \geq \frac{(q^{2n-2}-1)(q^{2n}-1)}{(q^2-1)(q^4-1)} > q^{4n-8}.$$

Finally, if $3 \leq k \leq n-3$, in particular $n \geq 6$, then

$$\chi(1) \geq (\widetilde{G} : C)_{p'}/2 \geq \frac{(q^{2n-4})(q^{2n-2}-1)(q^{2n}-1)}{2(q^2-1)(q^4-1)(q^6-1)} > q^{4n-8}.$$

□

6. Even characteristic orthogonal groups in even dimension

In this section we classify low-dimensional complex characters of the group $G = Spin^{\alpha}_{2n}(q) = \Omega^{\alpha}_{2n}(q)$, where $\alpha = \pm$ and q is a power of 2. Since q is even, we can identify G with its dual group. Note that, $\Omega^{\alpha}_{2n}(q) = [O^{\alpha}_{2n}(q), O^{\alpha}_{2n}(q)]$ and $O^{\alpha}_{2n}(q) = \Omega^{\alpha}_{2n}(q) \cdot \mathbb{Z}_2$.

Proof of Theorem 1.3. If χ is unipotent then we are done by Propositions 3.3, 3.4. Now we assume χ is not unipotent. Suppose that χ is parameterized by $((s), \psi)$, where s is a non-trivial semi-simple element in G and ψ is a unipotent character of $C := C_G(s)$ such that $\chi(1) \leq q^{4n-10}$. Set $C' = C_{O^{\alpha}_{2n}(q)}(s)$. Since C is a subgroup of C' of index 1 or 2, $(G : C)_{2'} = (O^{\alpha}_{2n}(q) : C')_{2'}$.

By Lemma 2.3, $C' \simeq O^{\pm}_{2m}(q) \times \prod_{i=1}^{t} GL^{\alpha_i}_{a_i}(q^{k_i})$, where $\alpha_i = \pm$, $\Sigma_{i=1}^{t} k_i a_i = n - m$. Since (s) is non-trivial, $m < n$. It is easy to see that $|\prod_{i=1}^{t} GL^{\alpha_i}_{a_i}(q^{k_i})|_{2'} \leq \prod_{i=1}^{n-m}(q^i - (-1)^i)$. Therefore, with convention that $q^0 - 1 = 1$, we have

$$(G : C)_{2'} \geq \frac{(q^{2(m+1)}-1)\cdots(q^{2n}-1)(q^m-1)}{(q+1)(q^2-1)\cdots(q^{n-m}-(-1)^{n-m})(q^n+1)} =: f(m,n).$$

1) When $1 \leq m \leq n-2$. We have

$$f(1,n) = (q-1)(q^n-1)\prod_{i=1}^{n-1}(q^i + (-1)^i) > q^{4n-10}$$

and

$$f(n-2,n) = \frac{(q^{2n-2}-1)(q^n-1)(q^{n-2}-1)}{(q+1)(q^2-1)} > q^{4n-10}.$$

Therefore, $\chi(1) \geq \min\{f(1,n), f(n-2,n)\} > q^{4n-10}$.

2) When $m = n-1$. Then $C \simeq O^{\pm}_{2n-2}(q) \times GL^{\beta}_1(q)$ with $\beta = \pm$. Since $GL^{\beta}_1(q) \simeq \Omega^{\beta}_2(q)$, $C' \simeq O^{\alpha\beta}_{2n-2}(q) \times GL^{\beta}_1(q)$. It is obvious that $C = \Omega^{\alpha\beta}_{2n-2}(q) \times GL^{\beta}_1(q)$. There are two cases:

- $C' = O^{\alpha}_{2n-2}(q) \times GL_1(q)$ corresponding to the case $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$, where $1 \neq \lambda \in \mathbb{F}^{\times}_q$. Note that there are $(q-2)/2$ choices for λ, namely, $\lambda = \gamma^k, k \in T_1$. For each such λ, there is exactly one semi-simple conjugacy class of such elements s in $O^{\alpha}_{2n}(q)$ by [W1, (3.7)]. Since $(G : C) = (O^{\alpha}_{2n}(q) : C')$, $O^{\alpha}_{2n}(q)$-conjugacy class of s is also a G-conjugacy of s. Therefore, there is also exactly one conjugacy class of semi-simple elements (s) in G such that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ for each $\lambda = \gamma^k, k \in T_1$.
- $C' = O^{-\alpha}_{2n-2}(q) \times GU_1(q)$ corresponding to the case $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$, where $1 \neq \lambda \in \mathbb{F}^{\times}_{q^2}$ and $\lambda^{-1} = \lambda^q$. Similarly as above, there is exactly one conjugacy class of semi-simple elements s in G such that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ for each $\lambda = \eta^k, k \in T_2$.

We have $\chi(1) = \frac{(q^n-\alpha)(q^{n-1}+\alpha\beta)}{q-\beta}\psi(1)$, where ψ is a unipotent character of C. If ψ is non-trivial and $(n, q, \alpha) \neq (5, 2, -)$ then by Propositions 7.1 and 7.2 of [TZ1],

$$\psi(1) \geq \frac{(q^{n-1}-\alpha\beta)(q^{n-2}+\alpha\beta q)}{q^2-1}.$$

In that case,

$$\chi(1) \geq \frac{(q^n-\alpha)(q^{n-1}+\alpha\beta)}{q-\beta} \cdot \frac{(q^{n-1}-\alpha\beta)(q^{n-2}+\alpha\beta q)}{q^2-1} > q^{4n-10}.$$

If ψ is non-trivial and $(n, q, \alpha) = (5, 2, -)$, then $\psi(1) \geq q^3(q-1)^3(q^3-1)/2 = 28$ and we still have $\chi(1) > q^{4n-10}$. So ψ must be trivial. In summary, in this case, χ is one of $(q-2)/2$ characters of degree $(q^n-\alpha)(q^{n-1}+\alpha)/(q-1)$ or $q/2$ characters of degree $(q^n-\alpha)(q^{n-1}-\alpha)/(q+1)$.

3) When $m = 0$. If $n \geq 6$, we still have $f(0, n) = \prod_{i=1}^{n}(q^i+(-1)^i)/(q^n+1) > q^{4n-10}$. Now we consider the case $n = 5$. Since $\chi(1) \leq q^{4n-10}$, $C' = GU_5(q)$. This forces $G = \Omega^-_{10}(q)$. Then $(G : C)_{2'} = (q-1)(q^2+1)(q^3-1)(q^4+1)$ and therefore $\psi(1) = 1$. There are exactly $q/2$ conjugacy classes (s) in $O^-_{10}(q)$ of semi-simple elements such that $C' \simeq GU_5(q)$, which is happened when $\mathrm{Spec}(s) = \{\lambda, \lambda, \lambda, \lambda, \lambda, \lambda^{-1}, \lambda^{-1}, \lambda^{-1}, \lambda^{-1}, \lambda^{-1}\}$, $\lambda = \eta^k, k \in T_2$. Note that $GU_5(q) \leq \Omega^-_{10}(q)$, so $C = GU_5(q)$ and $(O^-_{10}(q) : C') = 2(\Omega^-_{10}(q) : C)$. In other words, $|s^{O^-_{10}(q)}| = 2|s^{\Omega^-_{10}(q)}|$. Therefore there are exactly q conjugacy classes (s) in $\Omega^-_{10}(q)$ such that $C = GU_5(q)$. This gives q characters of degree $(q-1)(q^2+1)(q^3-1)(q^4+1)$ of $\Omega^-_{10}(q)$. $\square$

7. Odd characteristic orthogonal groups in even dimension

In this section, we classify low-dimensional complex representations of the group $G = Spin^{\alpha}_{2n}(q)$ where $\alpha = \pm$ and q is an odd prime power. The dual group G^* is the projective conformal orthogonal group $P(CO^{\alpha}_{2n}(q)^0)$, which is quotient of $CO^{\alpha}_{2n}(q)^0$ by its center $\mathbb{Z}(CO^{\alpha}_{2n}(q)^0) \simeq \mathbb{Z}_{q-1}$.

Proof of Theorem 1.4. If χ is unipotent then we are done by Propositions 3.3, 3.4. So we can assume that χ is not unipotent. We denote $\widetilde{G} := CO^{\alpha}_{2n}(q)$, $\widetilde{G}^0 := CO^{\alpha}_{2n}(q)^0$, and $Z := Z(\widetilde{G}) \simeq Z(\widetilde{G}^0) \simeq \mathbb{Z}_{q-1}$. Suppose that s^* is a non-trivial semi-simple element in G^*. Consider an inverse image s of s^* in $\widetilde{G}^0$. Then (s) is a non-trivial conjugacy class of semi-simple elements in $\widetilde{G}$. Set $C^* := C_{G^*}(s^*)$, $\widetilde{C} := C_{\widetilde{G}}(s)$, and $\widetilde{C}^0 := C_{\widetilde{G}^0}(s)$. Suppose that χ is parameterized by $((s^*), \psi), \psi \in \mathrm{Irr}(C^*)$, such that $\chi(1) \leq q^{4n-10}$. Let g be any element in $\widetilde{G}^0$ such that the image of g in G^* belongs to C^*. Then $gsg^{-1} = \mu(g)s$ for some $\mu(g) \in \mathbb{Z}_{q-1}$. Therefore, $\tau(s) = \tau(gsg^{-1}) = \tau(\mu(g)s) = \mu(g)^2\tau(s)$. It follows that $\mu(g)$ is 1 or -1. Hence,

$$(G^* : C^*)_{p'} \geq (\widetilde{G}^0 : \widetilde{C}^0)_{p'}/2 \geq (\widetilde{G} : \widetilde{C})_{p'}/4.$$

Case 1: If $\tau(s)$ is not a square in $\mathbb{F}_q$, multiplying s by a suitable scalar in $\mathbb{F}_q$, we can assume that $\tau(s)$ is fixed. By Lemma 2.5, $\widetilde{C} \simeq (O^{\pm}_m(q^2) \times \prod_{i=1}^t GL^{\alpha_i}_{a_i}(q^{k_i})) \cdot \mathbb{Z}_{q-1}$, where $\alpha_i = \pm$, $\Sigma_{i=1}^t k_i a_i = n - m$. Denote the group $\prod_{i=1}^t GL^{\alpha_i}_{a_i}(q^{k_i})$ by A for short. Now we will show that, when $n \geq 6$, $(\widetilde{G} : \widetilde{C})_{p'} > 4q^{4n-10}$, and therefore $\chi(1) \geq (G^* : C^*)_{p'} \geq (\widetilde{G} : \widetilde{C})_{p'}/4 > q^{4n-10}$. It is easy to see that $|A|_{p'} \leq \prod_{i=1}^{n-m}(q^i - (-1)^i)$. Hence,

$$(\widetilde{G} : \widetilde{C})_{p'} \geq \frac{2(q^2-1)\cdots(q^{2n}-1)}{(q^n+1)(q+1)(q^2-1)\cdots(q^{n-m}-(-1)^{n-m})|O^{\pm}_m(q^2|} =: f(m,n).$$

We have $f(0,n) = 2\prod_{i=1}^n(q^i + (-1)^i)/(q^n+1) > 4q^{4n-10}$ and $f(n,n) > 4q^{4n-10}$ for every $n \geq 6$. Therefore, $\chi(1) \geq (\widetilde{G} : \widetilde{C})_{p'}/4 \geq \min\{f(0,n), f(n,n)\}/4 > q^{4n-10}$ when $n \geq 6$.

Now let us consider the case $(n, \alpha) = (5, +)$. Note that $GU_5(q)$ is not a subgroup of $O^+_{10}(q)$. Therefore, if $A \neq GL_5(q)$, one can show that $(\widetilde{G} : \widetilde{C})_{p'} > 4q^{10}$ and hence $\chi(1) > q^{10}$. If $A = GL_5(q)$, then the characteristic polynomial of s has the form $P(x) = (x - \lambda)^5(x - \tau\lambda^{-1})^5$ with $\lambda \in \mathbb{F}^*_q$ and $\tau := \tau(s)$. Now we will show that $(G^* : C^*)_{p'} = (\widetilde{G}^0 : \widetilde{C}^0)_{p'}$ and therefore $\chi(1) \geq (\widetilde{G} : \widetilde{C})_{p'}/2 = (q+1)(q^2+1)(q^3+1)(q^4+1) > q^{10}$. Suppose that $s_1, s_2 \in \widetilde{G}^0$ are $\widetilde{G}^0$-conjugate and their images in G^* are the same. Then $s_1 = \pm s_2$. Let $V_1 := \mathrm{Ker}(s - \lambda)$ and $V_2 := \mathrm{Ker}(s - \tau\lambda^{-1})$. Note that V_1 and V_2 are totally isotropic subspaces in V and $V_1 \cap V_2 = \{0\}$. By definition, an element in $\widetilde{G}^0$ cannot carry V_1 to V_2. This implies that $s_1 = s_2$. In other words, $|s^{*G^*}| = |s^{\widetilde{G}^0}|$ and hence $(G^* : C^*)_{p'} = (\widetilde{G}^0 : \widetilde{C}^0)_{p'}$ as desired.

The next case is $(n, \alpha) = (5, -)$. Note that $GL_5(q)$ is not a subgroup of $O^-_{10}(q)$. Therefore, if $A \neq GU_5(q)$, one again can show that $\chi(1) > q^{10}$. If $A = GU_5(q)$, then the characteristic polynomial of s has the form $P(x) = [(x - \lambda)(x - \tau\lambda^{-1})]^5$, where $(x - \lambda)(x - \tau\lambda^{-1})$ is an irreducible polynomial over $\mathbb{F}_q$. Note that there are

$(q+1)/2$ choices for such a pair $(\lambda, \tau\lambda^{-1})$. So there are exactly $(q+1)/2$ conjugacy classes (s) in $\widetilde{G}$ such that $\widetilde{C} \simeq GU_5(q)$. That means there are exactly $(q+1)$ such conjugacy classes (s) in $\widetilde{G}^0$ since $\widetilde{C}^0 = \widetilde{C}$ in this case. Now modulo Z, we have exactly $(q+1)/2$ conjugacy classes (s^*) in G^*. Arguing similarly as above, one has $(G^* : C^*)_{p'} = (\widetilde{G}^0 : \widetilde{C}^0)_{p'} = (q-1)(q^2+1)(q^3-1)(q^4+1)$ and therefore $C^* = \widetilde{C}^0/Z$. The condition $\chi(1) = (G^* : C^*)_{p'}\psi(1) \le q^{10}$ implies that $\psi(1) = 1$. Since C^* has a unique unipotent character of degree 1, we get exactly $(q+1)/2$ characters of degree $(q-1)(q^2+1)(q^3-1)(q^4+1)$ in this case.

Case 2: If $\tau(s)$ is a square in $\mathbb{F}_q$, again, we can assume that $\tau(s) = 1$. By Lemma 2.5, $\widetilde{C} \simeq (O^{\pm}_{2k}(q) \times O^{\pm}_{2m-2k}(q) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i})) \cdot \mathbb{Z}_{q-1}$, where $\alpha_i = \pm$, $\Sigma_{i=1}^{t} k_i a_i = n-m$.

It is easy to see that $|O^{\pm}_{2k}(q) \times O^{\pm}_{2m-2k}(q)|_{p'} \le 2\prod_{i=1}^{m}(q^{2i}-1)/(q^m-1)$ if $m \ge 1$ and $|\prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(q^{k_i}|_{p'} \le \prod_{i=1}^{n-m}(q^i - (-1)^i)$. Therefore, with convention that $q^0 - 1 = 2$, we have

$$(\widetilde{G} : \widetilde{C})_{p'} \ge \frac{(q^{2(m+1)}-1)\cdots(q^{2n}-1)(q^m-1)}{(q^n+1)(q+1)(q^2-1)\cdots(q^{n-m}-(-1)^{n-m})} =: g(m,n).$$

1) When $1 \le m \le n-2$, $(\widetilde{G} : C)_{p'} \ge \min\{g(1,n), g(n-2,n)\}$. Direct computation shows that $g(1,n) = (q^n-1)\prod_{i=1}^{n-1}(q^i + (-1)^i)/(q+1) > 4q^{4n-10})$ and $g(n-2,n) = (q^{2n-2}-1)(q^n-1)(q^{n-2}-1)/(q+1)(q^2-1) > 4q^{4n-10}$. Therefore, $\chi(1) > q^{4n-10}$.

2) When $m = 0$, then $g(0,n) = 2\prod_{i=1}^{n}(q^i + (-1)^i)/(q^n+1)$. We still have $g(0,n) > 4q^{4n-10}$ if $n \ge 6$. The case $(n,\alpha) = (5,+)$ can be argued similarly as when $\tau(s)$ is not a square in $\mathbb{F}_q$. So we only need to consider the case $(n,\alpha) = (5,-)$. Note that $GL_5(q)$ is not a subgroup of $O^{-}_{10}(q)$. Therefore, if $A \ne GU_5(q)$, one again can show that $\chi(1) > q^{10}$. So we can assume $A = GU_5(q)$. Then the characteristic polynomial of s has the form $P(x) = [(x-\lambda)(x-\lambda^{-1})]^5$, where $(x-\lambda)(x-\lambda^{-1})$ is an irreducible polynomial over $\mathbb{F}_q$. Note that there are exactly $(q-1)/2$ such a pair (λ, λ^{-1}). Repeating arguments as when $\tau(s)$ is not a square in $\mathbb{F}_q$, we get exactly $(q-1)/2$ characters of degree $(q-1)(q^2+1)(q^3-1)(q^4+1)$ in this case.

3) When $m = n-1$. If $1 \le k \le n-2$ then $(\widetilde{G} : \widetilde{C})_{p'} \ge (q^{2n-2}-1)(q^n-1)(q^{n-2}-1)/2(q+1)^2 > 4q^{4n-10}$. Hence, $\chi(1) \ge (G^* : C^*)_{p'} > q^{4n-10}$. So $k = 0$ or $n-1$. With no loss, we can assume $k = n-1$. Note that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ in this case. If $\{1, ..., 1, \lambda, \lambda^{-1}\} = \{\mu, ..., \mu, \mu\lambda, \mu\lambda^{-1}\}$ for some $\mu \in \mathbb{F}_q^{\times}$, then $\mu = 1$ since $n \ge 5$. Therefore, $\widetilde{C}^0$ is the complete inverse image of C^* in $\widetilde{G}^0$. In other words, $C^* = \widetilde{C}^0/Z$ and hence $(G^* : C^*)_{p'} = (\widetilde{G}^0 : \widetilde{C}^0)_{p'} = (\widetilde{G} : \widetilde{C})_{p'}$. Also, since $GL_1^{\beta}(q) \simeq \Omega_2^{\beta}(q)$, $\widetilde{C} \simeq (O^{\alpha\beta}_{2n-2}(q) \times GL_1^{\beta}(q)) \cdot \mathbb{Z}_{q-1}$. One can show that $\widetilde{C}^0 \simeq (SO^{\alpha\beta}_{2n-2}(q) \times GL_1^{\beta}(q)) \cdot \mathbb{Z}_{q-1}$. There are two cases:

- $\widetilde{C} = (O^{\alpha}_{2n-2}(q) \times GL_1(q)) \cdot \mathbb{Z}_{q-1}$ corresponding to the case $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$, where $\pm 1 \ne \lambda \in \mathbb{F}_q^{\times}$. Note that there are $(q-3)/2$ choices for λ, namely, $\lambda = \gamma^k, k \in T_1$. For each such λ, there is exactly one semi-simple conjugacy class of such elements s in $O_{2n}^{\alpha}(q)$ by [W1, (2.6)]. This conjugacy class is also the conjugacy class of s in $\widetilde{G}^0$. Therefore, there is exactly one conjugacy class of semi-simple elements (s^*) in G^* such that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ for each $\lambda = \gamma^k, k \in T_1$.

- $\widetilde{C} = (O_{2n-2}^{-\alpha}(q) \times GU_1(q)) \cdot \mathbb{Z}_{q-1}$ corresponding to the case $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$, where $\pm 1 \neq \lambda \in \mathbb{F}_{q^2}^{\times}$ and $\lambda^{-1} = \lambda^q$. Note that there are $(q-1)/2$ choices for λ, namely, $\lambda = \eta^k, k \in T_2$. Similarly as above, there is exactly one conjugacy class of semi-simple elements (s^*) in G^* such that $\mathrm{Spec}(s) = \{1, ..., 1, \lambda, \lambda^{-1}\}$ for each $\lambda = \eta^k, k \in T_2$.

Using Proposition 2.1 for the canonical homomorphism $f : SO_{2n-2}^{\alpha\beta}(q) \times GL_1^{\beta}(q) \hookrightarrow \widetilde{C}^0 \to C^*$, $\beta = \pm$, we see that the unipotent characters of $SO_{2n-2}^{\alpha\beta}(q) \times GL_1^{\beta}(q)$ are of the form $\psi \circ f$, where ψ runs over the unipotent characters of C^*. In particular, by Propositions 7.1, 7.2 of [TZ1], ψ is trivial or $\psi(1) \geq (q^{n-1} - \alpha\beta)(q^{n-2} + \alpha\beta q)/(q^2 - 1)$. In the latter case,

$$\chi(1) \geq \frac{(q^n - \alpha)(q^{n-1} + \alpha\beta)}{q - \beta} \cdot \frac{(q^{n-1} - \alpha\beta)(q^{n-2} + \alpha\beta q)}{q^2 - 1} > q^{4n-10}.$$

Therefore, in this case, χ is one of $(q-3)/2$ representations of degree $(q^n - \alpha)(q^{n-1} + \alpha)/(q-1)$ or $(q-1)/2$ representations of degree $(q^n - \alpha)(q^{n-1} - \alpha)/(q+1)$.

4) When $m = n$. Since (s^*) is non-trivial, we assume $1 \leq k \leq n-1$.

If $k = 1$ or $n-1$, modulo Z, we may assume $\mathrm{Spec}(s) = \{-1, -1, 1, ..., 1\}$. Then $\widetilde{C} = (O_2^{\beta}(q) \times O_{2n-2}^{\alpha\beta}(q)) \cdot \mathbb{Z}_{q-1}$ and $\widetilde{C}^0 \simeq (SO_2^{\beta}(q) \times SO_{2n-2}^{\alpha\beta}(q)) \cdot 2 \cdot \mathbb{Z}_{q-1}$. Again, $C^* = \widetilde{C}^0/Z$ and $(G^* : C^*)_{p'} = (\widetilde{G} : \widetilde{C})_{p'}$. There is a unique possibility for (s^*) for each $\beta = \pm$. We have $\chi(1) = \frac{(q^n-\alpha)(q^{n-1}+\alpha\beta)}{2(q-\beta)}\psi(1)$ where ψ is a unipotent character of C^*. Consider the canonical homomorphism $f : (SO_2^{\beta}(q) \times SO_{2n-2}^{\alpha\beta}(q)) \cdot 2 \hookrightarrow \widetilde{C}^0 \to C^*$, whose kernel is contained in the center of $(SO_2^{\beta}(q) \times SO_{2n-2}^{\alpha\beta}(q)) \cdot 2$ and image contains the commutator group of C^* since $(SO_2^{\beta}(q) \times SO_{2n-2}^{\alpha\beta}(q)) \cdot 2$ contains the commutator group of $\widetilde{C}^0$. By Proposition 2.1, the unipotent characters of $(SO_2^{\beta}(q) \times SO_{2n-2}^{\alpha\beta}(q)) \cdot 2$ are of the form $\psi \circ f$, where ψ runs over the unipotent characters of C^*. In particular, ψ is one of two linear unipotent characters or $\psi(1) \geq (q^{n-1} - \alpha\beta)(q^{n-2} + \alpha\beta q)/(q^2 - 1)$. In the latter case,

$$\chi(1) \geq \frac{(q^n - \alpha)(q^{n-1} + \alpha\beta)}{2(q - \beta)} \cdot \frac{(q^{n-1} - \alpha\beta)(q^{n-2} + \alpha\beta q)}{q^2 - 1} > q^{4n-10}.$$

If $2 \leq k \leq n-2$, then

$$\chi(1) \geq \frac{(\widetilde{G} : \widetilde{C})_{p'}}{4} \geq \frac{|O_{2n}^{\alpha}(q)|}{4|O_4^{\beta}(q)| \cdot |O_{2n-4}^{\alpha\beta}(q)|} =$$

$$= \frac{(q^{2n-2}-1)(q^n-\alpha)(q^{n-2}+\alpha\beta)(q^2+\beta)}{8(q^2-1)(q^4-1)} > q^{4n-10}.$$

□

8. Groups $Spin_{12}^{\pm}(3)$

In this section, we will classify the irreducible complex characters of $G = Spin_{12}^{\alpha}(3)$, $\alpha = \pm$, of degrees up to $4 \cdot 3^{15}$. Note that $G^* = P(CO_{12}^{\alpha}(3)^0)$. The argument in this section is similar to that in §7. We will keep all notation from there. Two following lemmas can be checked by direct computation.

Lemma 8.1. *Suppose that* $\chi \in \mathrm{Irr}(P(CO_{12}^{-}(3)^0))$ *is unipotent. Then either* χ *is one of the characters labeled by* $\binom{0\ 6}{-}$, $\binom{1\ 5}{-}$, $\binom{2\ 4}{-}$, $\binom{0\ 1\ 6}{1}$, $\binom{0\ 1\ 2}{5}$, *or* $\chi(1) > 4 \cdot 3^{15}$.

Lemma 8.2. *Suppose that* $\chi \in \mathrm{Irr}(P(CO_{12}^{+}(3)^0))$ *is unipotent. Then either* χ *is one of characters labeled by* $\binom{6}{0}$, $\binom{5}{1}$, $\binom{4}{2}$, $\binom{3}{3}$ *(2 characters),* $\binom{0\ 1}{1\ 6}$, $\binom{0\ 1\ 2\ 5}{-}$, *or* $\chi(1) > 4 \cdot 3^{15}$.

Proposition 8.3. $Spin_{12}^{+}(3)$ *has exactly* 28 *irreducible complex characters of degrees less than* $4 \cdot 3^{15}$ *and* $Spin_{12}^{-}(3)$ *has exactly* 16 *irreducible complex characters of degrees less than* $4 \cdot 3^{15}$.

Proof. Let χ be an irreducible complex character of G of degree less than $4 \cdot 3^{15}$. Since we have already counted the number of unipotent characters in two above lemmas, we can assume that χ is not unipotent. Suppose that χ is parameterized by $((s^*), \psi)$, where $1 \neq s^* \in G^* := P(CO_{12}^{\alpha}(3)^0)$ and $\psi \in \mathrm{Irr}(C^*)$, $C^* = C_{G^*}(s^*)$. Recall that $\widetilde{G} := CO_{12}^{\alpha}(3)$ and $\widetilde{G}^0 := CO_{12}^{\alpha}(3)^0$. Denote $Z := Z(\widetilde{G}) = Z(\widetilde{G}^0)$ the center of $\widetilde{G}$ as well as $\widetilde{G}^0$. Let s be an inverse image of s^* in $\widetilde{G}^0$. Set $\widetilde{C} = C_{\widetilde{G}}(s)$ and $\widetilde{C}^0 = C_{\widetilde{G}^0}(s)$.

Case 1: $\tau(s) = -1$. Then we have $\widetilde{C} \simeq (O_m^{\pm}(9) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(3^{k_i})) \cdot \mathbb{Z}_2$, where $\alpha_i = \pm$, $\Sigma_{i=1}^{t} k_i a_i + m = 6$. Since $\chi(1) < 4 \cdot 3^{15}$, $(\widetilde{G} : \widetilde{C})_{3'} < 16 \cdot 3^{15}$. This inequality happens only when $\widetilde{C} \simeq GL_6^{\pm}(3) \cdot \mathbb{Z}_2$. This forces $G = Spin_{12}^{+}(3)$. Let us consider the case $\widetilde{C} \simeq GL_6(3) \cdot \mathbb{Z}_2$. Note that there is a unique conjugacy class (s) of semi-simple elements in $\widetilde{G}$ so that $\widetilde{C} \simeq GL_6(3) \cdot \mathbb{Z}_2$, which is happened when the characteristic polynomial of s is $(x^2-1)^6$. In this case, $\widetilde{C}^0 = \widetilde{C}$, $(\widetilde{G} : \widetilde{C}) = 2(\widetilde{G}^0 : \widetilde{C}^0)$, and therefore there are two conjugacy classes of semi-simple elements in $\widetilde{G}^0$, as well as in G^*, such that $\widetilde{C} \simeq GL_6(3) \cdot \mathbb{Z}_2$. Furthermore, C^* is an extension by 2 of $\widetilde{C}/Z$. Therefore, χ is one of 4 characters of degree $(G^* : C^*)_{3'} = \frac{1}{2}\prod_{i=1}^{5}(3^i+1)$, provided that $G = Spin_{12}^{+}(3)$.

Next, we consider the case $\widetilde{C} \simeq GU_6(3) \cdot \mathbb{Z}_2$. Note that there are two semi-simple conjugacy classes (s) in $\widetilde{G}$ so that $\widetilde{C} \simeq GL_6(3) \cdot \mathbb{Z}_2$, which is happened when the characteristic polynomial of s is $(x^2+x-1)^6$ or $(x^2-x-1)^6$. In this case, $\widetilde{C}^0 = \widetilde{C}$,

$(\widetilde{G} : \widetilde{C}) = 2(\widetilde{G}^0 : \widetilde{C}^0)$, and therefore there are 4 such conjugacy classes of semi-simple elements in $\widetilde{G}^0$. Modulo Z, we get exactly two semi-simple conjugacy classes (s^*) in G^*. Furthermore, $C^* \simeq \widetilde{C}/Z$. Therefore, we get two characters of degree $\prod_{i=1}^{5}(3^i + (-1)^i)$ of $G = Spin_{12}^+(3)$.

Case 2: $\tau(s) = 1$. We have $\widetilde{C} \simeq (O_{2k}^{\pm}(3) \times O_{2m-2k}^{\pm}(3) \times \prod_{i=1}^{t} GL_{a_i}^{\alpha_i}(3^{k_i})) \cdot \mathbb{Z}_2$, where $\alpha_i = \pm$, $\Sigma_{i=1}^{t} k_i a_i + m = 6$. The inequality $\chi(1) < 4 \cdot 3^{15}$ implies that $m = 0, 5$, or 6. If $m = 0$, similarly as in case 1, one can show that $\widetilde{C} \simeq GU_6(3) \cdot \mathbb{Z}_2$ and C^* is an extension by 2 of $\widetilde{C}/Z$. Hence, χ is one of 4 characters of degree $\frac{1}{2}\prod_{i=1}^{5}(3^i + (-1)^i)$, provided that $G = Spin_{12}^+(3)$.

When $m = 5$, it is easy to show that $(G^* : C^*)_{3'} = (\widetilde{G} : \widetilde{C})_{3'}$. If $k = 0$ or 5, we have already shown in the proof of Theorem 1.4 that either χ is the unique character of degree $(3^6 - \alpha)(3^5 - \alpha)/4$ or

$$\chi(1) \geq \frac{(3^6 - \alpha)(3^5 - \alpha)}{4} \cdot \frac{(3^5 + \alpha)(3^4 - \alpha 3)}{8} > 4 \cdot 3^{15}.$$

If $1 \leq k \leq 4$, $\chi(1) \geq (\widetilde{G} : \widetilde{C})_{3'} \geq (3^{10} - 1)(3^6 - 1)(3^4 - 1)/32 > 4 \cdot 3^{15}$.

Now we suppose $m = 6$. First, If $k = 1$ or 5, we have $\chi = \psi(1)(3^6 - \alpha)(3^5 + \alpha\beta)/2(3 - \beta)$, where $\beta = \pm$ and ψ is a unipotent character of $C^* = \widetilde{C}^0/Z$, $\widetilde{C}^0 \simeq (SO_2^{\beta}(3) \times SO_{10}^{\alpha\beta}(3)) \cdot 2 \cdot \mathbb{Z}_2$. When $\beta = +$, the inequality $\chi(1) < 4 \cdot 3^{15}$ forces ψ to be linear. Therefore, χ is one of 2 characters of degree $(3^6 - \alpha)(3^5 + \alpha)/4$. When $\beta = -$, note that the unipotent characters of $(SO_2^-(3) \times SO_{10}^{-\alpha}(3)) \cdot 2$ are of the form $\psi \circ f$, where f is the canonical homomorphism $f : (SO_2^-(3) \times SO_{10}^{-\alpha}(3)) \cdot 2 \to C^*$. Since $SO_2^-(3)$ has a unique unipotent character which is trivial, the unipotent characters of $(SO_2^-(3) \times SO_{10}^{-\alpha}(3)) \cdot 2$ are actually unipotent characters of $O_{10}^{-\alpha}(3)$. Therefore, ψ is one of 2 unipotent characters of degree 1; 2 unipotent characters of degree $(3^5 + \alpha)(3^4 - \alpha 3)/8$; or $\psi(1) \geq (3^{10} - 3^2)/8$. Hence there are 4 characters of degrees less than $4 \cdot 3^{15}$ in this case. Next, if $k = 2$ or 4, we have

$$\chi(1) = \frac{(3^4 + \alpha\beta)(3^{10} - 1)(3^6 - \alpha)}{2(3^2 - 1)(3^2 - \beta)} \cdot \psi(1),$$

where ψ is a unipotent character of $C^* = \widetilde{C}^0/Z(\widetilde{G})$, $\widetilde{C}^0 \simeq (SO_4^{\beta}(3) \times SO_8^{\alpha\beta}(3)) \cdot 2 \cdot \mathbb{Z}_2$. Since $\chi(1) < 4 \cdot 3^{15}$, ψ is one of two linear unipotent characters of C^* for each $\beta = \pm$. This gives 4 more characters of degrees less than $4 \cdot 3^{15}$. Finally, if $k = 3$, then $\chi(1) \geq (G^* : C^*)_{3'} \geq (\widetilde{G} : \widetilde{C})_{3'}/2 > 4 \cdot 3^{15}$. $\square$

References

[BK] J. Brundan and A. S. Kleshchev, Lower bounds for the degrees of irreducible Brauer characters of finite general linear groups, *J. Algebra* **223** (2000), 615 – 629.

[C1] R. W. Carter, *'Finite groups of Lie type. Conjugacy classes and complex characters'*, Wiley and Sons, New York et al, 1985, 544 pp.

[C2] R. W. Carter, Centralizers of semisimple elements in the finite classical groups, *Proc. London Math. Soc.* **42** (1981), 1 – 41.

[CH] M. Claßen-Houben, Die unipotenten Charaktere fur die GAP-Charaktertafeln der endlichen Gruppen vom Lie-Typ, Diplomarbeit, RWTH Aachen, 2005.

[DM] F. Digne and J. Michel, *'Representations of finite groups of Lie type'*, London Mathematical Society Student Texts 21, 1991, 159 pp.

[FS] P. Fong and B. Srinivasan, The blocks of finite classical groups, *J. Reine Angew. Math.* **396** (1989), 122 – 191.

[GMST] R. M. Guralnick, K. Magaard, J. Saxl and P. H. Tiep, Cross characteristic representations of symplectic and unitary groups, *J. Algebra* **257** (2002), 291 – 347.

[GT1] R. M. Guralnick and P. H. Tiep, Low-dimensional representations of special linear groups in cross characteristic, *Proc. London Math. Soc.* **78** (1999), 116 – 138.

[GT2] R. M. Guralnick and P. H. Tiep, Cross characteristic representations of even characteristic symplectic groups, *Trans. Amer. Math. Soc.* **356** (2004), 4969 – 5023.

[HM] G. Hiss and G. Malle, Low dimensional representations of special unitary groups, *J. Algebra* **236** (2001), 745 – 767.

[LOST] M. W. Liebeck, E. A. O'Brien, A. Shalev, and P. H. Tiep, The Ore conjecture, submitted.

[LS] V. Landazuri and G. Seitz, On the minimal degrees of projective representations of the finite Chevalley groups, *J. Algebra* **32** (1974), 418 – 443.

[Lu] F. Lübeck, Smallest degrees of representations of exceptional groups of Lie type, *Comm. Algebra* **29** (2001), 2147 – 2169.

[SZ] G. Seitz and A. E. Zalesskii, On the minimal degrees of projective representations of the finite Chevalley groups, II, *J. Algebra* **158** (1993), 233 – 243.

[ST] P. Sin and P. H. Tiep, Rank 3 permutation modules of the finite classical groups, *J. Algebra* **291** (2005), 551 – 606.

[S] B. Srinivasan, The characters of the finite symplectic group $Sp(4, q)$, *Trans. Amer. Math. Soc.* **131** (1968), 488 – 525.

[T] P. H. Tiep, Low dimensional representations of finite quasisimple groups, *Groups, combinatorics and geometry* (Durham, 2001), World Sci. Publishing, River Edge, NJ, (2003), 277–294.

[TZ1] P. H. Tiep and A. E. Zalesskii, Minimal characters of the finite classical groups, *Comm. Algebra* **24** (1996), 2093 – 2167.

[TZ2] P. H. Tiep and A. E. Zalesskii, Unipotent elements of finite groups of Lie type and realization fields of their complex representations, *J. Algebra* **271** (2004), 327 – 390.

[W1] G. E. Wall, On the conjugacy classes in the unitary, symplectic and orthogonal groups, *J. Austral. Math. Soc.* **3** (1963), 1 – 62.

[W2] D. White, The 2-decomposition numbers of $Sp(4, q)$, q odd, *J. Algebra* **131** (1990), 703 – 725.

Department of Mathematics, Michigan State University, East Lansing, MI 48824, USA

E-mail address: `hungnguyen@math.msu.edu`